



The Ehrenfeucht-Fraïssé-Game of Length #1

Alan Mekler; Saharon Shelah; Jouko Väänänen

Transactions of the American Mathematical Society, Vol. 339, No. 2. (Oct., 1993), pp. 567-580.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28199310%29339%3A2%3C567%3ATEOL%3E2.0.CO%3B2-8>

Transactions of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

THE EHRENFUCHT-FRAÏSSÉ-GAME OF LENGTH ω_1

ALAN MEKLER, SAHARON SHELAH, AND JOUKO VÄÄNÄNEN

ABSTRACT. Let $\mathfrak{A}$ and $\mathfrak{B}$ be two first order structures of the same vocabulary. We shall consider the *Ehrenfeucht-Fraïssé-game of length ω_1 of $\mathfrak{A}$ and $\mathfrak{B}$* which we denote by $\mathcal{E}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$. This game is like the ordinary Ehrenfeucht-Fraïssé-game of $L_{\omega\omega}$ except that there are ω_1 moves. It is clear that $\mathcal{E}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined if $\mathfrak{A}$ and $\mathfrak{B}$ are of cardinality $\leq \aleph_1$. We prove the following results:

Theorem 1. *If $V = L$, then there are models $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\aleph_2$ such that the game $\mathcal{E}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is nondetermined.*

Theorem 2. *If it is consistent that there is a measurable cardinal, then it is consistent that $\mathcal{E}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\leq \aleph_2$.*

Theorem 3. *For any $\kappa \geq \aleph_3$ there are $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality κ such that the game $\mathcal{E}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is nondetermined.*

1. INTRODUCTION

Let $\mathfrak{A}$ and $\mathfrak{B}$ be two first order structures of the same vocabulary L . We denote the domains of $\mathfrak{A}$ and $\mathfrak{B}$ by A and B respectively. All vocabularies are assumed to be relational.

The *Ehrenfeucht-Fraïssé-game of length γ of $\mathfrak{A}$ and $\mathfrak{B}$* , denoted by $\mathcal{E}_\gamma(\mathfrak{A}, \mathfrak{B})$, is defined as follows: There are two players called $\forall$ and $\exists$. First $\forall$ plays x_0 and then $\exists$ plays y_0 . After this $\forall$ plays x_1 , and $\exists$ plays y_1 , and so on. If $\langle (x_\beta, y_\beta) : \beta < \alpha \rangle$ has been played and $\alpha < \gamma$, then $\forall$ plays x_α after which $\exists$ plays y_α . Eventually a sequence $\langle (x_\beta, y_\beta) : \beta < \gamma \rangle$ has been played. The rules of the game say that both players have to play elements of $A \cup B$. Moreover, if $\forall$ plays his x_β in A (B), then $\exists$ has to play his y_β in B (A). Thus the sequence $\langle (x_\beta, y_\beta) : \beta < \gamma \rangle$ determines a relation $\pi \subseteq A \times B$. Player $\exists$ wins this round of the game if π is a partial isomorphism. Otherwise $\forall$ wins. The notion of winning strategy is defined in the usual manner. We say that a player *wins $\mathcal{E}_\gamma(\mathfrak{A}, \mathfrak{B})$* if he has a winning strategy in $\mathcal{E}_\gamma(\mathfrak{A}, \mathfrak{B})$.

Received by the editors May 31, 1991.

1991 *Mathematics Subject Classification.* Primary 03C75; Secondary 03C55.

The first author's research partially supported by NSERC grant #A8948.

The second author's research partially supported by the United States Israel Binational Science Foundation; Publication number 416.

The third author's research partially supported by the Academy of Finland.

Recall that

$$\begin{aligned}\mathfrak{A} \equiv_{\omega\omega} \mathfrak{B} &\Leftrightarrow \forall n < \omega (\exists \text{ wins } \mathcal{G}_n(\mathfrak{A}, \mathfrak{B})) \\ \mathfrak{A} \equiv_{\infty\omega} \mathfrak{B} &\Leftrightarrow \exists \text{ wins } \mathcal{G}_\omega(\mathfrak{A}, \mathfrak{B}).\end{aligned}$$

In particular, $\mathcal{G}_\gamma(\mathfrak{A}, \mathfrak{B})$ is determined for $\gamma \leq \omega$. The question, whether $\mathcal{G}_\gamma(\mathfrak{A}, \mathfrak{B})$ is determined for $\gamma > \omega$, is the subject of this paper. We shall concentrate on the case $\gamma = \omega_1$.

The notion

$$(1) \quad \exists \text{ wins } \mathcal{G}_\gamma(\mathfrak{A}, \mathfrak{B})$$

can be viewed as a natural generalization of $\mathfrak{A} \equiv_{\infty\omega} \mathfrak{B}$. The latter implies isomorphism for countable models. Likewise (1) implies isomorphism for models of cardinality $|\gamma|$:

Proposition 1. *Suppose $\mathfrak{A}$ and $\mathfrak{B}$ have cardinality $\leq \kappa$. Then $\mathcal{G}_\kappa(\mathfrak{A}, \mathfrak{B})$ is determined: $\exists$ wins if $\mathfrak{A} \cong \mathfrak{B}$, and $\forall$ wins if $\mathfrak{A} \not\cong \mathfrak{B}$.*

Proof. If $f: \mathfrak{A} \cong \mathfrak{B}$, then the winning strategy of $\exists$ in $\mathcal{G}_\kappa(\mathfrak{A}, \mathfrak{B})$ is to play in such a way that the resulting π satisfies $\pi \subseteq f$. On the other hand, if $\mathfrak{A} \not\cong \mathfrak{B}$, then the winning strategy of $\forall$ is to systematically enumerate $A \cup B$ so that the final π will satisfy $A = \text{dom}(\pi)$ and $B = \text{rng}(\pi)$. $\square$

For models of arbitrary cardinality we have the following simple but useful criterion of (1), namely in the terminology of [15] that they are “potentially isomorphic.” We use $\text{Col}(\lambda, \kappa)$ to denote the notion of forcing which collapses $|\lambda|$ to κ (with conditions of cardinality less than κ).

Proposition 2. *Suppose $\mathfrak{A}$ and $\mathfrak{B}$ have cardinality $\leq \lambda$ and κ is regular. Player $\exists$ wins $\mathcal{G}_\kappa(\mathfrak{A}, \mathfrak{B})$ if and only if $\Vdash_{\text{Col}(\lambda, \kappa)} \mathfrak{A} \cong \mathfrak{B}$.*

Proof. Suppose τ is a winning strategy of $\exists$ in $\mathcal{G}_\kappa(\mathfrak{A}, \mathfrak{B})$. Since $\text{Col}(\lambda, \kappa)$ is $< \kappa$ -closed,

$$\Vdash_{\text{Col}(\lambda, \kappa)} \text{“}\tau \text{ is a winning strategy of } \exists \text{ in } G_\kappa(\mathfrak{A}, \mathfrak{B}).\text{”}$$

Hence $\Vdash_{\text{Col}(\lambda, \kappa)} \mathfrak{A} \cong \mathfrak{B}$ by Proposition 1. Suppose then $p \Vdash \tilde{f}: \mathfrak{A} \cong \mathfrak{B}$ for some $p \in \text{Col}(\lambda, \kappa)$. While the game $\mathcal{G}_\kappa(\mathfrak{A}, \mathfrak{B})$ is played, $\exists$ keeps extending the condition p further and further. Suppose he has extended p to q and $\forall$ has played $x \in A$. Then $\exists$ finds $r \leq q$ and $y \in B$ with $r \Vdash \tilde{f}(x) = y$. Using this simple strategy $\exists$ wins. $\square$

Proposition 3. *Suppose T is an ω -stable first order theory with NDOP. Then $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all models $\mathfrak{A}$ of T and all models $\mathfrak{B}$.*

Proof. Suppose $\mathfrak{A}$ is a model of T . If $\mathfrak{B}$ is not $L_{\infty\omega_1}$ -equivalent to $\mathfrak{A}$, then $\forall$ wins $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ easily. So let us suppose $\mathfrak{A} \equiv_{\infty\omega_1} \mathfrak{B}$. We may assume A and B are of cardinality $\geq \aleph_1$. If we collapse $|A|$ and $|B|$ to $\aleph_1$, T will remain ω -stable with NDOP, and $\mathfrak{A}$ and $\mathfrak{B}$ will remain $L_{\infty\omega_1}$ -equivalent. So $\mathfrak{A}$ and $\mathfrak{B}$ become isomorphic by [19, Chapter XIII, §1]. Now Proposition 2 implies that $\exists$ wins $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$. $\square$

Hyttinen [10] showed that $\mathcal{G}_\gamma(\mathfrak{A}, \mathfrak{B})$ may be nondetermined for all γ with $\omega < \gamma < \omega_1$ and asked whether $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ may be nondetermined. Our results

show that $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ may be nondetermined for $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\aleph_3$ (Theorem 17), but for models of cardinality $\aleph_2$ the answer is more complicated.

Let $F(\omega_1)$ be the free group of cardinality $\aleph_1$. Using the combinatorial principle $\square_{\omega_1}$ we construct an abelian group G of cardinality $\aleph_2$ such that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is nondetermined (Theorem 4). On the other hand, we show that starting with a model with a measurable cardinal one can build a forcing extension in which $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all models $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\leq \aleph_2$ (Theorem 14).

Thus the free abelian group $F(\omega_1)$ has the remarkable property that the question

Is $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ determined for all G ?

cannot be answered in ZFC alone. Proposition 3 shows that no model of an $\aleph_1$ -categorical first order theory can have this property.

We follow Jech [11] in set theoretic notation. We use S_n^m to denote the set $\{\alpha < \omega_m : \text{cf}(\alpha) = \omega_n\}$. Closed and unbounded sets are called cub sets. A set of ordinals is λ -closed if it is closed under supremums of ascending λ -sequences $\langle \alpha_i : i < \lambda \rangle$ of its elements. A subset of a cardinal is λ -stationary if it meets every λ -closed unbounded subset of the cardinal. The closure of a set A of ordinals in the order topology of ordinals is denoted by $\overline{A}$. The free abelian group of cardinality κ is denoted by $F(\kappa)$.

2. A NONDETERMINED $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ WITH G A GROUP OF CARDINALITY $\aleph_2$

In this section we use $\square_{\omega_1}$ to construct a group G of cardinality $\aleph_2$ such that the game $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is nondetermined (Theorem 4). For background on almost free groups the reader is referred to [4]. However, our presentation does not depend on special knowledge of almost free groups. All groups below are assumed to be abelian.

By $\square_{\omega_1}$ we mean the principle, which says that there is a sequence $\langle C_\alpha : \alpha < \omega_2, \alpha = \bigcup \alpha \rangle$ such that

1. C_α is a cub subset of α .
2. If $\text{cf}(\alpha) = \omega$, then $|C_\alpha| = \omega$.
3. If γ is a limit point of C_α , then $C_\gamma = C_\alpha \cap \gamma$.

Recall that $\square_{\omega_1}$ follows from $V = L$ by a result of R. Jensen [14]. For a sequence of sets C_α as above we can let

$$E_\beta = \{\alpha \in S_0^2 : \text{the order type of } C_\alpha \text{ is } \beta\}.$$

For some $\beta < \omega_1$ the set E_β has to be stationary. Let us use E to denote this E_β . Then E is a so-called *nonreflecting* stationary set, i.e., if $\text{cf}(\gamma) > \omega$ then $E \cap \gamma$ is nonstationary on γ . Indeed, then some final segment D_γ of the set of limit points of C_γ is a cub subset of γ disjoint from E . Moreover, $\text{cf}(\alpha) = \omega$ for all $\alpha \in E$.

Theorem 4. *Assuming $\square_{\omega_1}$, there is a group G of cardinality $\aleph_2$ such that the game $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is nondetermined.*

Proof. Let $\mathbb{Z}^{\omega_2}$ denote the direct product of ω_2 copies of the additive group $\mathbb{Z}$ of the integers. Let x_α be the element of $\mathbb{Z}^{\omega_2}$ which is 0 on coordinates $\neq \alpha$ and 1 on the coordinate α . Let us fix for each $\delta \in S_0^2$ an ascending cofinal

sequence $\eta_\delta: \omega \rightarrow \delta$. For such δ , let

$$z_\delta = \sum_{n=0}^{\infty} 2^n x_{\eta_\delta(n)}.$$

Let $\langle C_\alpha: \alpha = \bigcup \alpha < \omega_2 \rangle$, $\langle D_\alpha: \alpha = \bigcup \alpha < \omega_2 \rangle$, and $E = E_\beta$ be obtained from $\square_{\omega_1}$ as above. We are ready to define the groups we need for the proof: Let G be the smallest pure subgroup of $\mathbb{Z}^{\omega_2}$ which contains x_α for $\alpha < \omega_2$ and z_δ for $\delta \in E$, let G_α be the smallest pure subgroup of $\mathbb{Z}^{\omega_2}$ which contains x_γ for $\gamma < \alpha$ and z_δ for $\delta \in E \cap \alpha$, let $F (= F(\omega_2))$ be the subgroup of $\mathbb{Z}^{\omega_2}$ generated freely by x_α for $\alpha < \omega_2$, and finally, let F_α be the subgroup of $\mathbb{Z}^{\omega_2}$ generated freely by x_γ for $\gamma < \alpha$.

The properties we shall want of G_α are standard but for the sake of completeness we shall sketch proofs. We need that each G_α is free and for any $\beta \notin E$ any free basis of G_β can be extended to a free basis of G_α for all $\alpha > \beta$.

The proof is by induction on α . For limit ordinals we use the fact that E is nonreflecting. The case of successors of ordinals not in E is also easy. Assume now that $\delta \in E$ and the induction hypothesis has been verified up to δ . By the induction hypothesis for any $\beta < \delta$ such that $\beta \notin E$, there is n_0 so that $G_\delta = G_\beta \oplus H \oplus K$ where K is the group freely generated by $\{x_{\eta_\delta(n)}: n_0 \leq n\}$ and $x_{\eta_\delta(m)} \in G_\beta$ for all $m < n_0$. Then $G_{\delta+1} = G_\beta \oplus H \oplus K'$ where K' is freely generated by $\{\sum_{m=n}^{\infty} 2^{m-n} x_{\eta_\delta(m)}: n_0 \leq n\}$.

On the other hand, if $\delta \in E$ and $\{x_{\eta_\delta(n)}: n < \omega\} \subseteq B$, where B is a subgroup of G such that $z_\delta \notin B$, then G/B is nonfree, as $z_\delta + B$ is infinitely divisible by 2 in G/B .

Claim 1. $\exists$ does not win $\mathcal{G}_{\omega_1}(F, G)$.

Suppose τ is a winning strategy of $\exists$. Let $\alpha \in E$ such that the pair (G_α, F_α) is closed under the first ω moves of τ , that is, if $\forall$ plays his first ω moves inside $G_\alpha \cup F_\alpha$, then τ orders $\exists$ to do the same. We shall play $\mathcal{G}_{\omega_1}(F, A)$ pointing out the moves of $\forall$ and letting τ determine the moves of $\exists$. On his move number $2n$ $\forall$ plays the element $x_{\eta_\alpha(n)}$ of G_α . On his move number $2n+1$ $\forall$ plays some element of F_α . Player $\forall$ plays his moves in F_α in such a way that during the first ω moves eventually some countable direct summand K of F_α as well as some countable $B \subseteq G_\alpha$ are enumerated. Let J be the smallest pure subgroup of G containing $B \cup \{z_\alpha\}$. During the next ω moves of $\mathcal{G}_{\omega_1}(F, A)$ player $\forall$ enumerates J and $\exists$ responds by enumerating some $H \subseteq F$. Since τ is a winning strategy, H has to be a subgroup of F . But now H/K is free, whereas J/B is nonfree, so $\forall$ will win the game, a contradiction.

Claim 2. $\forall$ does not win $\mathcal{G}_{\omega_1}(F, G)$.

Suppose τ is a winning strategy of $\forall$. If we were willing to use CH, we could just take α of cofinality ω_1 such that (F_α, G_α) is closed under τ , and derive a contradiction from the fact that $F_\alpha \cong G_\alpha$. However, since we do not want to assume CH, we have to appeal to a longer argument.

Let $\kappa = (2^\omega)^{++}$. Let $\mathfrak{M}$ be the expansion of $\langle H(\kappa), \in \rangle$ obtained by adding the following structure to it:

- (H1) The function $\delta \mapsto \eta_\delta$.
- (H2) The function $\delta \mapsto z_\delta$.

- (H3) The function $\alpha \mapsto C_\alpha$.
 (H4) A well-ordering $<$ of the universe.
 (H5) The winning strategy τ .

Let $\mathfrak{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of $\mathfrak{M}$ such that $\omega_1 \subseteq N$ and $N \cap \omega_2$ is an ordinal α of cofinality ω_1 .

Let $D_\alpha = \{\beta_i : i < \omega_1\}$ in ascending order. Since $C_{\beta_i} = C_\alpha \cap \beta_i$, every initial segment of C_α is in N . By elementarity, $G_{\beta_i} \in N$ for all $i < \omega_1$. Let ϕ be an isomorphism $G_\alpha \rightarrow F_\alpha$ obtained as follows: ϕ restricted to G_{β_0} is the $<$ -least isomorphism between the free groups G_{β_0} and F_0 . If ϕ is defined on all G_{β_j} , $j < i$, then ϕ is defined on G_{β_i} as the $<$ -least extension of $\bigcup_{j < i} \phi_{\beta_j}$ to an isomorphism between G_{β_i} and F_i . Recall that by our choice of D_α , $G_{\beta_{i+1}}/G_{\beta_i}$ is free, so such extensions really exist.

We derive a contradiction by showing that $\exists$ can play ϕ against τ for the whole duration of the game $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$. To achieve this we have to show that when $\exists$ plays his canonical strategy based on ϕ , the strategy τ of $\forall$ directs $\forall$ to go on playing elements which are in N , that is, elements of $G_\alpha \cup F_\alpha$.

Suppose a sequence $s = \langle (x_\gamma, y_\gamma) : \gamma < \mu \rangle$, $\mu < \omega_1$, has been played. It suffices to show that $s \in N$. Choose β_i so that the elements of s are in $G_{\beta_i} \cup F_{\beta_i}$. Now s is uniquely determined by $\phi \upharpoonright G_{\beta_i}$ and τ . Note that because $C_{\beta_i} = C_\alpha \cap \beta_i$, $\phi \upharpoonright G_{\beta_i}$ can be defined inside N similarly as ϕ was defined above, using C_{β_i} instead of C_α . Thus $s \in N$ and we are done.

We have proved that $\mathcal{G}_{\omega_1}(F, G)$ is nondetermined. This clearly implies $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is nondetermined. $\square$

Remark. R. Jensen [14, p. 286] showed that if $\square_{\omega_1}$ fails, then ω_2 is Mahlo in L . Therefore, if $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all almost free groups $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\aleph_2$, then ω_2 is Mahlo in L . If we start with $\square_\kappa$, we get an almost free group A of cardinality κ^+ such that $\mathcal{G}_{\omega_1}(F(\omega_1), A)$ is nondetermined.

3. $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ CAN BE DETERMINED FOR ALL G

In this section all groups are assumed to be abelian. It is easy to see that $\exists$ wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ for any uncountable free group G , so in this exposition $F(\omega_1)$ is a suitable representative of all free groups. In the study of determinacy of $\mathcal{G}_{\omega_1}(F(\omega_1), \mathfrak{A})$ for various $\mathfrak{A}$ it suffices to study groups $\mathfrak{A}$, since for other $\mathfrak{A}$ player $\forall$ easily wins the game.

Starting from a model with a Mahlo cardinal we construct a forcing extension in which $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined, when G is any group of cardinality $\aleph_2$. This can be extended to groups G of any cardinality, if we start with a supercompact cardinal.

In the proof of the next results we shall make use of *stationary logic* $L(aa)$ introduced in [17]. For the definition and basic facts about $L(aa)$ the reader is referred to [1]. This logic has a new quantifier $aa s$ quantifying over variables s ranging over countable subsets of the universe. A cub set of such s is any set which contains a superset of any countable subset of the universe and which is closed under unions of countable chains. The semantics of $aa s$ is defined as follows:

$$aa s \phi(s, \dots) \Leftrightarrow \phi(s, \dots) \text{ holds for a cub set of } s.$$

Note that a group of cardinality $\aleph_1$ is free if and only if it satisfies

(+) $aa \ s \ aa \ s' (s \subseteq s' \rightarrow s \text{ is free and } s'/s \text{ is free}).$

Proposition 5. *Let G be a group of cardinality at most $\aleph_2$. Then the following conditions are equivalent:*

- (1) $\exists$ wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$.
- (2) G satisfies (+).
- (3) G is the union of a continuous chain $\langle G_\alpha : \alpha < \omega_2 \rangle$ of free subgroups with $G_{\alpha+1}/G_\alpha$ $\aleph_1$ -free for all $\alpha < \omega_2$.

Proof. (1) implies (2): Suppose $\exists$ wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$. By Proposition 2 we have $\Vdash_{\text{Col}(|G|, \omega_1)}$ “ G is free”. Using the countable completeness of $\text{Col}(|G|, \omega_1)$ it is now easy to construct a cub set S of countable subgroups of G such that if $A \in S$ then for all $B \in S$ with $A \subseteq B$ we have B/A free. Thus G satisfies (+). (2) implies (3) quite trivially. (3) implies (1): Suppose a continuous chain as in (3) exists. If we collapse $|G|$ to $\aleph_1$, then in the extension the chain has length $< \omega_2$. Now we use Theorem 1 of [8]:

If a group A is the union of a continuous chain of $< \omega_2$ free subgroups $\{A_\alpha : \alpha < \gamma\}$ of cardinality $\leq \aleph_1$ such that each $A_{\alpha+1}/A_\alpha$ is $\aleph_1$ -free, then A is free.

Thus G is free in the extension and (1) follows from Proposition 2. $\square$

Remark. Conditions (1) and (2) of Proposition 5 are equivalent for G of any size.

Let us consider the following principle:

- (*) For all stationary $E \subseteq S_0^2$ and countable subsets a_α of $\alpha \in E$ such that a_α is cofinal in α and of order type ω there is a closed $C \subseteq \omega_2$ of order type ω_1 such that $\{\alpha \in E : a_\alpha \setminus C \text{ is finite}\}$ is stationary in C .

Lemma 6. *The principle (*) implies that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G of cardinality $\aleph_2$.*

Proof. Suppose G is a group of cardinality $\aleph_2$. We may assume the domain of G is ω_2 . Let us assume G is $\aleph_2$ -free, as otherwise $\forall$ easily wins. We prove that G satisfies condition (3) of Proposition 5 and thereby that $\exists$ wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$.

To prove condition (3) of Proposition 5, assume the contrary. Then G can be expressed as the union of a continuous chain $\langle G_\alpha : \alpha < \omega_2 \rangle$ of free groups with $G_{\alpha+1}/G_\alpha$ non- $\aleph_1$ -free for $\alpha \in E$, $E \subseteq \omega_2$ stationary. By Fodor’s Lemma, we may assume $E \subseteq S_0^2$ or $E \subseteq S_1^2$. The latter case is much easier and therefore we assume $E \subseteq S_0^2$. Also we may assume that for all α , every ordinal in $G_{\alpha+1} \setminus G_\alpha$ is greater than every ordinal in G_α . Finally by intersecting with a closed unbounded set we may assume that for all $\alpha \in E$ the set underlying G_α is α . Choose for each $\alpha \in E$ some countable subgroup b_α of $G_{\alpha+1}$ with $b_\alpha + G_\alpha/G_\alpha$ nonfree. Let $c_\alpha = b_\alpha \cap G_\alpha$. We will choose a_α so that any final segment generates a subgroup containing c_α . Enumerate c_α as $\{g_n : n < \omega\}$ such that each element is enumerated infinitely often. Choose an increasing sequence $(\alpha_n : n < \omega)$ cofinal in α so that for all n , $g_n \in G_{\alpha_n}$. Finally, for each n , choose $h_n \in G_{\alpha_{n+1}} \setminus G_{\alpha_n}$. Let $a_\alpha = \{h_n : n < \omega\} \cup \{h_n + g_n : n < \omega\}$. It

is now easy to check that a_α is a sequence of order type ω which is cofinal in α and any subgroup of G which contains all but finitely many of the elements of a_α contains c_α .

By (*) there is a continuous C of order type ω_1 such that $\{\alpha \in C: a_\alpha \setminus C \text{ is finite}\}$ is stationary in C . Let $D = \langle C \cup \sum_{\alpha \in C} b_\alpha \rangle$. Since $|D| \leq \aleph_1$, D is free.

For any $\alpha \in C$, let

$$D_\alpha = \left\langle (C \cap \alpha) \cup \left(\sum_{\beta \in (C \cap \alpha)} b_\beta \right) \right\rangle.$$

Note that $D = \bigcup_{\alpha \in C} D_\alpha$, each D_α is countable and for limit point δ of C , $D_\delta = \bigcup_{\alpha \in (C \cap \delta)} D_\alpha$. Hence there is an $\alpha \in C \cap E$ such that $a_\alpha \setminus C$ is finite and D/D_α is free. Hence $b_\alpha + D_\alpha/D_\alpha$ is free. But $b_\alpha + D_\alpha/D_\alpha \cong b_\alpha/b_\alpha \cap D_\alpha = b_\alpha/b_\alpha \cap G_\alpha$, which is not free, a contradiction. $\square$

For the next theorem we need a lemma from [6]. A proof is included for the convenience of the reader.

Lemma 7 [6]. *Suppose λ is a regular cardinal and $\mathbb{Q}$ is a notion of forcing which satisfies the λ -c.c. Suppose $\mathcal{I}$ is a normal λ -complete ideal on λ and $\mathcal{I}^+ = \{S \subseteq \lambda: S \notin \mathcal{I}\}$. For all sets $S \in \mathcal{I}^+$ and sequences of conditions $\langle p_\alpha: \alpha \in S \rangle$, there is a set C with $\lambda \setminus C \in \mathcal{I}$ so that for all $\alpha \in C \cap S$,*

$$p_\alpha \Vdash_{\mathbb{Q}} \{ \beta: p_\beta \in \tilde{G} \} \in \mathcal{I}^+, \text{ where } \mathcal{I} \text{ is the ideal generated by } \mathcal{I}^+.$$

Proof. Suppose the lemma is false. So there is an $\mathcal{I}$ -positive set $S' \subseteq S$ such that for all $\alpha \in S'$ there is an extension r_α of p_α and a set $I_\alpha \in \mathcal{I}$ (note: I_α is in the ground model) so that

$$r_\alpha \Vdash \{ \beta: p_\beta \in \tilde{G} \} \subseteq I_\alpha.$$

Let I be the diagonal union of $\{I_\alpha: \alpha \in S'\}$.

Suppose now that $\alpha < \beta$ and $\alpha, \beta \in (S' \setminus I)$. Since $\beta \notin I$, $r_\alpha \Vdash p_\beta \notin \tilde{G}$. Hence $r_\alpha \Vdash r_\beta \notin \tilde{G}$. So r_α, r_β are incompatible. Hence $\{r_\alpha: \alpha \in S' \setminus I\}$ is an antichain which, since S' is $\mathcal{I}$ -positive, is of cardinality λ . This is a contradiction. $\square$

Remark. It is a well-known fact that the ideal $\mathcal{I}$ of Lemma 7 is forced by $\mathbb{Q}$ to be λ -complete and normal.

Theorem 8. *Assuming the consistency of a Mahlo cardinal, it is consistent that (*) holds and hence that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G of cardinality $\aleph_2$.*

Proof. By a result of Harrington and Shelah [7] we may start with a Mahlo cardinal κ in which every stationary set of cofinality ω reflects, that is, if $S \subseteq \kappa$ is stationary, and $\text{cf}(\alpha) = \omega$ for $\alpha \in S$, then $S \cap \lambda$ is stationary in λ for some inaccessible $\lambda < \kappa$.

For any inaccessible λ let $\mathbb{P}_\lambda$ be the Levy-forcing for collapsing λ to ω_2 . The conditions of $\mathbb{P}_\lambda$ are countable functions $f: \lambda \times \omega_1 \rightarrow \lambda$ such that $f(\alpha, \beta) < \alpha$ for all α and β and each f is increasing and continuous in the second coordinate. It is well known that $\mathbb{P}_\lambda$ is countably closed and satisfies the λ -chain condition [11, p. 191].

Let $\mathbb{P} = \mathbb{P}_\kappa$. Suppose $p \in \mathbb{P}$ and

$$p \Vdash \text{“}\tilde{E} \subseteq S_0^2 \text{ is stationary and} \\ \forall \alpha \in \tilde{E} (\tilde{a}_\alpha \subseteq \alpha \text{ is cofinal in } \alpha \text{ and of order type } \omega)\text{.”}$$

Let

$$S = \{\alpha < \kappa : \exists q \leq p (q \Vdash \alpha \in \tilde{E})\}.$$

For any $\alpha \in S$ let $p_\alpha \leq p$ such that $p_\alpha \Vdash \alpha \in \tilde{E}$. Since $\mathbb{P}$ is countably closed, we can additionally require that for some countable $a_\alpha \subseteq \alpha$ we have $p_\alpha \Vdash \tilde{a}_\alpha = a_\alpha$.

The set S is stationary in κ , for if $C \subseteq \kappa$ is cub, then $p \Vdash C \cap \tilde{E} \neq \emptyset$, whence $C \cap S \neq \emptyset$. Also $\text{cf}(\alpha) = \omega$ for $\alpha \in S$. Let λ be inaccessible such that $S \cap \lambda$ is stationary in λ . We may choose λ in such a way that $\alpha \in S \cap \lambda$ implies $p_\alpha \in \mathbb{P}_\lambda$. By Lemma 7 there is a $\delta \in S \cap \lambda$ such that

$$p_\delta \Vdash_{\mathbb{P}_\lambda} \text{“}\tilde{E}_1 = \{\alpha < \lambda : p_\alpha \in \tilde{G}\} \text{ is stationary.”}$$

Let $\mathbb{Q}$ be the set of conditions $f \in \mathbb{P}$ with $\text{dom}(f) \subseteq (\kappa \setminus \lambda) \times \omega_1$. Note that $\mathbb{P} \cong \mathbb{P}_\lambda \otimes \mathbb{Q}$. Let G be $\mathbb{P}$ -generic containing p_δ and $G_\lambda = G \cap \mathbb{P}_\lambda$ for any inaccessible $\lambda \leq \kappa$. Then G_κ is $\mathbb{P}_\lambda$ -generic and ω_2 of $V[G_\lambda]$ is λ . Let us work now in $V[G_\lambda]$. Thus λ is the current ω_2 , $E_1 = \{\alpha < \lambda : p_\alpha \in G_\lambda\}$ is stationary, and we have the countable sets $a_\alpha \subseteq \alpha$ for $\alpha \in E_1$. Since $\mathbb{Q}$ collapses λ there is a name $\tilde{f}$ such that

$$\Vdash_{\mathbb{Q}} \text{“}\tilde{f} : \omega_1 \rightarrow \lambda \text{ is continuous and cofinal.”}$$

More precisely $\tilde{f}$ is the name for the function f defined by $f(\alpha) = \beta$ if and only if there is some $g \in G$ so that $g(\lambda, \alpha) = \beta$. Let $\tilde{C}$ denote the range of $\tilde{f}$. We shall prove the following statement:

Claim. $\Vdash_{\mathbb{Q}} \{\alpha < \lambda : a_\alpha \setminus \tilde{C} \text{ is finite}\}$ is stationary in $\tilde{C}$.

Suppose $q \in \mathbb{Q}$ so that $q \Vdash \text{“}\tilde{D} \subseteq \omega_1 \text{ is a cub.”}$ Let $\mathfrak{M}$ be an appropriate expansion of $\langle H(\kappa), \in \rangle$ and $\langle \mathfrak{N}_i : i < \lambda \rangle$, $\mathfrak{N}_i = \langle N_i, \in, \dots \rangle$, a sequence of elementary submodels of $\mathfrak{M}$ such that

- (i) Everything relevant is in N_0 .
- (ii) If $\alpha_i = N_i \cap \lambda$, then $\alpha_i < \alpha_j$ for $i < j < \lambda$.
- (iii) N_{i+1} is closed under countable sequences.
- (iv) $|N_i| = \omega_1$.
- (v) $N_i = \bigcup_{j < i} N_j$ for i a limit ordinal.

Choose $\gamma = \alpha_i \in E_1$ and let $\langle i_n : n < \omega \rangle$ be a sequence of successor ordinals such that $\gamma = \sup\{\alpha_{i_n} : n < \omega\}$. Let $q_0 \leq q$ and $\beta_0 \in \omega_1$ such that $q_0, \beta_0 \in N_{i_0}$,

$$q_0 \Vdash \text{“}\beta_0 \in \tilde{D}\text{”}$$

and q_0 decides the value of $\tilde{f}''\beta_0$ (which will by elementarity necessarily be a subset of α_{i_0}).

If q_n and β_n are defined we choose $q_{n+1} \leq q_n$ and $\beta_{n+1} \in \omega_1$ such that $q_{n+1}, \beta_{n+1} \in N_{i_{n+1}}$,

$$q_{n+1} \Vdash \text{“}\beta_{n+1} \in \tilde{D} \text{ and } a_\gamma \cap (\alpha_{i_{n+1}} \setminus \alpha_{i_n}) \subseteq \tilde{f}''\beta_{n+1} \subseteq \alpha_{i_{n+1}}\text{”}$$

and q_{n+1} decides $\tilde{f}''\beta_{n+1}$. Finally, let

$$q_\omega = \bigcup \{q_n : n < \omega\} \quad \text{and} \quad \beta = \bigcup \{\beta_n : n < \omega\}.$$

Then

$$q_\omega \Vdash \text{“}\beta \in \tilde{D} \quad \text{and} \quad a_\gamma \setminus \tilde{f}''\beta \text{ is finite.”}$$

The claim, and thereby the theorem, is proved. $\square$

Corollary 9. *The statement that $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for every structure $\mathfrak{A}$ of cardinality $\aleph_2$ and every uncountable free group $\mathfrak{B}$, is equiconsistent with the existence of a Mahlo cardinal.*

Remark. If $\mathcal{G}_{\omega_1}(A, F(\omega_1))$ is determined for all groups A of cardinality κ^+ , κ singular, then $\square_\kappa$ fails. This implies that the Covering Lemma fails for the Core Model, whence there is an inner model for a measurable cardinal. This shows that the conclusion of Theorem 8 cannot be strengthened to arbitrary G . However, by starting with a larger cardinal we can make this extension:

Theorem 10. *Assuming the consistency of a supercompact cardinal, it is consistent that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G .*

Proof. Let us assume that the stationary logic $L_{\omega_1\omega}(aa)$ has the Löwenheim-Skolem property down to $\aleph_1$. This assumption is consistent relative to the consistency of a supercompact cardinal [2]. Let G be an arbitrary $\aleph_2$ -free group. Let H be an $L(aa)$ -elementary submodel of G of cardinality $\aleph_1$. Thus H is a free group. The group H satisfies the sentence $(+)$, whence so does G . Now the claim follows from Proposition 5. $\square$

Corollary 11. *Assuming the consistency of a supercompact cardinal, it is consistent that $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for every structure $\mathfrak{A}$ and every uncountable free group $\mathfrak{B}$.*

4. $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ CAN BE DETERMINED FOR ALL $\mathfrak{A}$ AND $\mathfrak{B}$ OF CARDINALITY $\aleph_2$

We prove the consistency of the statement that $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\leq \aleph_2$ assuming the consistency of a measurable cardinal. Actually we make use of an assumption that we call $I^*(\omega)$ concerning stationary subsets of ω_2 . This assumption is known to imply that ω_2 is measurable in an inner model. It follows from the previous section that some large cardinal axioms are needed to prove the stated determinacy.

Let $I^*(\omega)$ be the following assumption about ω_1 -stationary subsets of ω_2 :

$I^*(\omega)$: Let $\mathcal{I}$ be the ω_1 -nonstationary ideal NS_{ω_1} on ω_2 . Then $\mathcal{I}^+$ has a σ -closed dense subset K .

Hodges and Shelah [9] define a principle $I(\omega)$, which is like $I^*(\omega)$ except that $\mathcal{I}$ is not assumed to be the ω_1 -nonstationary ideal. They use $I(\omega)$ to prove the determinacy of an Ehrenfeucht-Fraïssé-game played on several boards simultaneously.

Note that $I^*(\omega)$ implies $\mathcal{I}$ is precipitous, so the consistency of $I^*(\omega)$ implies the consistency of a measurable cardinal [12]. For the proof of the following result the reader is referred to [12]:

Theorem 12 (Laver). *The assumption $I^*(\omega)$ is consistent relative to the consistency of a measurable cardinal.*

We shall consider models $\mathfrak{A}$, $\mathfrak{B}$ of cardinality $\aleph_2$, so we may as well assume they have ω_2 as universe. For such $\mathfrak{A}$ and $\alpha < \omega_2$ we let $\mathfrak{A}_\alpha$ denote the structure $\mathfrak{A} \cap \alpha$. Similarly $\mathfrak{B}_\alpha$.

Lemma 13. *Suppose $\mathfrak{A}$ and $\mathfrak{B}$ are structures of cardinality $\aleph_2$. If $\forall$ does not have a winning strategy in $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$, then $S = \{\alpha : \mathfrak{A}_\alpha \cong \mathfrak{B}_\alpha\}$ is ω_1 -stationary.*

Proof. Let $C \subseteq \omega_2$ be ω_1 -closed and unbounded. Suppose $S \cap C = \emptyset$. We derive a contradiction by describing a winning strategy of $\forall$: Let $\pi : \omega_1 \times \omega_1 \times 2 \rightarrow \omega_1$ be onto with $\alpha, \beta, d \leq \pi(\alpha, \beta, d)$ for all $\alpha, \beta < \omega_1$ and $d < 2$. If $\alpha < \omega_2$, let $\theta_\alpha : \omega_1 \rightarrow \alpha$ be onto. Suppose the sequence $\langle (x_i, y_i) : i < \alpha \rangle$ has been played. Here x_i denotes a move of $\forall$ and y_i a move of $\exists$. During the game $\forall$ has built an ascending sequence $\{c_i : i < \alpha\}$ of elements of C . Now he lets c_α be the smallest element of C greater than all the elements x_i, y_i , $i < \alpha$. Suppose $\alpha = \pi(i, \gamma, d)$. Now $\forall$ will play $\theta_{c_i}(\gamma)$ as an element of $\mathfrak{A}$, if $d = 0$, and as an element of $\mathfrak{B}$ if $d = 1$.

After all ω_1 moves of $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ have been played, some $\mathfrak{A}_\alpha$ and $\mathfrak{B}_\alpha$, where $\alpha \in C$, have been enumerated. Since $\alpha \notin S$, $\forall$ has won the game. $\square$

Theorem 14. *Assume $I^*(\omega)$. The game $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is determined for all $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\leq \aleph_2$.*

Proof. Suppose $\forall$ does not have a winning strategy. By Lemma 13 the set $S = \{\alpha : \mathfrak{A}_\alpha \cong \mathfrak{B}_\alpha\}$ is ω_1 -stationary. Let I and K be as in $I^*(\omega)$. If $\alpha \in S$, let $h_\alpha : \mathfrak{A}_\alpha \cong \mathfrak{B}_\alpha$. We describe a winning strategy of $\exists$. The idea of this strategy is that $\exists$ lets the isomorphisms h_α determine his moves. Of course, different h_α may give different information to $\exists$, so he has to decide which h_α to follow. The key point is that $\exists$ lets some h_α determine his move only if there are stationarily many other h_β that agree with h_α on this move.

Suppose the sequence $\langle (x_i, y_i) : i < \alpha \rangle$ has been played. Again x_i denotes a move of $\forall$ and y_i a move of $\exists$. Suppose $\forall$ plays next x_α and this is (say) in A . During the game $\exists$ has built a descending sequence $\{S_i : i < \alpha\}$ of elements of K with $S_0 \subseteq S$. The point of the sets S_i is that $\exists$ has taken care that for all $i < \alpha$ and $\beta \in S_i$ we have $y_i = h_\beta(x_i)$ or $x_i = h_\beta(y_i)$ depending on whether $\forall$ played x_i in A or B . Now $\exists$ lets $S'_\alpha \subseteq \bigcap_{i < \alpha} S_i$ so that $S'_\alpha \in K$ and $\forall i \in S'_\alpha$ ($x_\alpha < i$). For each $i \in S'_\alpha$ we have $h_i(x_\alpha) < i$. By normality, there are an $S_\alpha \subseteq S'_\alpha$ in K and a y_α such that $h_i(x_\alpha) = y_\alpha$ for all $i \in S_\alpha$. This element y_α is the next move of $\exists$. Using this strategy $\exists$ wins. $\square$

5. A NONDETERMINED $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ WITH $\mathfrak{A}$ AND $\mathfrak{B}$ OF CARDINALITY $\aleph_3$

We construct directly in ZFC two models $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\aleph_3$ with $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ nondetermined. It readily follows that such models exist in all cardinalities $\geq \aleph_3$. The construction uses a square-like principle (Lemma 16), which is provable in ZFC.

Lemma 15 [18]. *There is a stationary $X \subseteq S_1^3$ and a sequence $\langle D_\alpha : \alpha \in X \rangle$ such that*

1. D_α is a cub subset of α for all $\alpha \in X$.

2. The order type of D_α is ω_1 .
3. If $\alpha, \beta \in X$ and $\gamma < \min\{\alpha, \beta\}$ is a limit of both D_α and D_β , then $D_\alpha \cap \gamma = D_\beta \cap \gamma$.
4. If $\gamma \in D_\alpha$, then γ is a limit point of D_α if and only if γ is a limit ordinal.

Proof. See [18, Theorem 4.1] or [3, Lemma 7.7]. $\square$

Lemma 16. *There are sets S, T , and C_α for $\alpha \in S$ such that the following hold:*

1. $S \subseteq S_0^3 \cup S_1^3$ and $S \cap S_1^3$ is stationary.
2. $T \subseteq S_0^3$ is stationary and $S \cap T = \emptyset$.
3. If $\alpha \in S$, then $C_\alpha \subseteq \alpha \cap S$ is closed and of order-type $\leq \omega_1$.
4. If $\alpha \in S$ and $\beta \in C_\alpha$, then $C_\beta = C_\alpha \cap \beta$.
5. If $\alpha \in S \cap S_1^3$, then C_α is cub on α .

Proof. Let X and $\langle D_\alpha : \alpha \in X \rangle$ be as in Lemma 15. Let $S' = X \cup Y$, where Y consists of ordinals which are limit points $< \alpha$ of some D_α , $\alpha \in X$. If $\alpha \in X$, we let C_α be the set of limit points $< \alpha$ of D_α . If $\alpha \in Y$, we let C_α be the set of limit points $< \alpha$ of $D_\beta \cap \alpha$, where $\beta > \alpha$ is chosen arbitrarily from X .

Now claims 1, 3, 4 and 6 are clearly satisfied.

Let $S_0^3 = \bigcup_{i < \omega_2} T_i$ where the T_i are disjoint stationary sets. Since $|\overline{C_\alpha}| \leq \omega_1$, there is $i_\alpha < \omega_2$ such that $i \geq i_\alpha$ implies $\overline{C_\alpha} \cap T_i = \emptyset$. Let $S'' \subseteq S'$ be stationary such that $\alpha \in S''$ implies i_α is constant i . Let $T = T_i$. Finally, let $S = S'' \cup \bigcup \{C_\alpha : \alpha \in S''\}$. Claim 2 is satisfied, and the lemma is proved. $\square$

Theorem 17. *There are structures $\mathfrak{A}$ and $\mathfrak{B}$ of cardinality $\aleph_3$ with one binary predicate such that the game $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is nondetermined.*

Proof. Let S, T , and $\langle C_\alpha : \alpha \in S \rangle$ be as in Lemma 16. We shall construct a sequence $\{M_\alpha : \alpha < \omega_3\}$ of sets and a sequence $\{G_\alpha : \alpha \in S\}$ of functions such that the conditions (M1)–(M5) below hold. Let W_α be the set of all mappings $G_{\gamma_0}^{d_0} \dots G_{\gamma_n}^{d_n}$, where $\gamma_0, \dots, \gamma_n \in S \cap \alpha$, $d_i \in \{-1, 1\}$, G_γ^1 means G_γ and G_γ^{-1} means the inverse of G_γ . Let $W = W_{\omega_3}$. (Note that W consists of a set of partial functions.)

The conditions on the M_α 's and the G_α 's are

(M1) $M_\alpha \subseteq M_\beta$ if $\alpha < \beta$, and $M_\alpha \subset M_{\alpha+1}$ if $\alpha \in S$.

(M2) $M_\nu = \bigcup_{\alpha < \nu} M_\alpha$ for limit ν .

(M3) G_α is a bijection of $M_{\alpha+1}$ for $\alpha \in S$.

(M4) If $\beta \in S$ and $\alpha \in C_\beta$, then $G_\alpha \subseteq G_\beta$.

(M5) If for some β , $G_\beta(a) = b$ and for some $w \in W$, $w(a) = b$, then there is some γ so that $w \subseteq G_\gamma$. Furthermore if β is the minimum ordinal so that $G_\beta(a) = b$ then $\gamma = \beta$ or $\beta \in C_\gamma$.

In order to construct the set $M = \bigcup_{\alpha < \omega_3} M_\alpha$ and the mappings G_α we define an oriented graph with M as the set of vertices. We use the terminology of Serre [16] for graph-theoretic notions. If x is an edge, the origin of x is denoted by $o(x)$ and the terminus by $t(x)$. Our graph has an inverse edge $\bar{x}$ for each edge x . Thus $o(\bar{x}) = t(x)$ and $t(\bar{x}) = o(x)$. Some edges are called *positive*, the rest are called *negative*. An edge is positive if and only if its inverse is negative. For each edge x of M there is a set $L(x)$ of labels. The set of possible labels for positive edges is $\{g_\alpha : \alpha < \omega_3\}$. The negative edges can have elements of $\{g_\alpha^{-1} : \alpha < \omega_3\}$ as labels. The labels are assumed to be given in such a way that a positive edge gets g_α as a label if and only if its inverse gets

the label g_α^{-1} . During the construction the sets of labels will be extended step by step.

The construction is analogous to building an acyclic graph on which a group acts freely. The graph then turns out to be the Cayley graph of the group. The labelled graph we will build will be the “Cayley graph” of W which will be as free as possible given (M1)–(M4). Condition (M5) is a consequence of the freeness of the construction.

Let us suppose the sets M_β , $\beta < \alpha$, of vertices have been defined. Let $M_{<\alpha} = \bigcup_{\beta < \alpha} M_\beta$. Some vertices in $M_{<\alpha}$ have edges between them and a set $L(x)$ of labels has been assigned to each such edge x .

If α is a limit ordinal, we let $M_\alpha = M_{<\alpha}$. So let us assume $\alpha = \beta + 1$. If $\beta \notin S$, $M_\alpha = M_\beta$. So let us assume $\beta \in S$. Let $\gamma = \sup(C_\beta)$. Notice that since S consists entirely of limit ordinals and $C_\beta \subseteq S$, either $\gamma = \beta$ or $\gamma + 1 < \beta$.

Case 1. $\gamma = \beta$: We extend M_β to M_α by adding new vertices $\{P_z : z \in \mathbb{Z}\}$ and for each $z \in \mathbb{Z}$ a positive edge $x_\alpha^{P_z}$ with $o(x_\alpha^{P_z}) = P_z$ and $t(x_\alpha^{P_z}) = P_{z+1}$. We also let $L(x_\alpha^{P_z}) = \{g_\beta\} \cup \{g_\delta : \beta \in C_\delta\}$.

Case 2. $\gamma + 1 < \beta$: We extend M_β to M_α by adding new vertices $\{P'_z : z \in \mathbb{Z} \setminus \{0\}\}$ for each $P \in M_\beta \setminus M_{\gamma+1}$. For notational convenience let $P'_0 = P$. Now we add for each $P \in M_\beta \setminus M_{\gamma+1}$ new edges as follows. For each $z \in \mathbb{Z}$ we add a positive edge $x_\alpha^{P'_z}$ with

$$o(x_\alpha^{P'_z}) = P'_z, \quad t(x_\alpha^{P'_z}) = P'_{z+1}, \quad L(x_\alpha^{P'_z}) = \{g_\beta\} \cup \{g_\delta : \beta \in C_\delta\}.$$

This determines completely the inverse of $x_\alpha^{P'_z}$.

This ends the construction of the graph. In the construction each vertex P in $M_{\alpha+1}$, $\alpha \in S$, is made the origin of a unique edge x_α^P with $g_\alpha \in L(x_\alpha^P)$. We define $G_\alpha(P) = t(x_\alpha^P)$.

The construction of the sets M_α and the mappings G_α is now completed. It follows immediately from the construction that each G_α , $\alpha \in S$, is a bijection of $M_{\alpha+1}$. So (M1)–(M3) hold. (M4) holds, because g_α is added to the labels of any edge with g_β , where $\beta \in C_\alpha$, as a label. Finally, (M5) is a consequence of the fact that the graph is circuit-free.

Let us fix $a_0 \in M_1$ and $b_0 = G_{\beta_0}(a_0)$, where $\beta_0 \in C_\alpha$ for all $\alpha \in S$. Note that we may assume, without loss of generality, the existence of such a β_0 .

If $a_0, a_1 \in M$, let

$$R_{(a_0, a_1)} = \{(a'_0, a'_1) \in M^2 : \exists w \in W(w(a_0) = a'_0 \wedge w(a_1) = a'_1)\}.$$

We let

$$\mathfrak{M} = \langle M, (R_{(a_0, a_1)})_{(a_0, a_1) \in M^2} \rangle, \quad \mathfrak{A} = \langle \mathfrak{M}, a_0 \rangle, \quad \mathfrak{B} = \langle \mathfrak{M}, b_0 \rangle,$$

and show that $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ is nondetermined.

The reduction of the language of $\mathfrak{A}$ and $\mathfrak{B}$ to one binary predicate is easy. One just adds a copy of ω_3 , together with its ordering, and a copy of $M \times M$ to the structures with the projection maps. Then fix a bijection ϕ from ω_3 to M^2 .

Add a new binary predicate R to the language and interpret R to be contained in $\omega_3 \times M^2$ such that $R(\beta, (a, b))$ holds if and only if $R_{\phi(\beta)}(a, b)$ holds. We can now dispense with the old binary predicates. We have replaced our structure by one in a finite language without making any difference to who wins the game $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$. The extra step of reducing to a single binary predicate is standard.

An important property of these models is that if $\alpha \in S \cap S_1^3$, then $G_\alpha \upharpoonright M_\alpha$ is an automorphism of the restriction of $\mathfrak{M}$ to M_α and takes a_0 to b_0 .

Claim 3. $\forall$ does not win $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$.

Suppose $\forall$ has a winning strategy τ . Again, there is a quick argument which uses CH: Find $\alpha \in S$ such that M_α is closed under τ and $\text{cf}(\alpha) = \omega_1$. Now C_α is cub on α , whence G_α maps M_α onto itself. Using G_α player $\exists$ can easily beat τ , a contradiction.

In the following longer argument we need not assume CH. Let κ be a large regular cardinal. Let $\mathfrak{H}$ be the expansion of $\langle H(\kappa), \in \rangle$ obtained by adding the following structure to it:

- (H1) The function $\alpha \mapsto M_\alpha$.
- (H2) The function $\alpha \mapsto G_\alpha$.
- (H3) The function $\alpha \mapsto C_\alpha$.
- (H4) A well-ordering $<^*$ of the universe.
- (H5) The winning strategy τ .
- (H6) The sets S and T .

Let $\mathfrak{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of $\mathfrak{H}$ of cardinality $\aleph_2$ such that $\alpha = N \cap \omega_3 \in S \cap S_1^3$.

Now C_α is a cub of order-type ω_1 on α and G_α maps M_α onto M_α . Moreover, G_α is a partial isomorphism from $\mathfrak{A}$ into $\mathfrak{B}$. Provided that τ does not lead $\forall$ to play his moves outside M_α , $\exists$ has an obvious strategy: he lets G_α determine his moves. So let us assume a sequence $\langle (x_\xi, y_\xi) : \xi < \gamma \rangle$ has been played inside M_α and $\gamma < \omega_1$. Let $\beta \in C_\alpha$ such that M_β contains the elements x_ξ, y_ξ for $\xi < \gamma$. The sequence $\langle y_\xi : \xi < \gamma \rangle$ is totally determined by G_β and τ . Since $G_\beta \in N$, $\langle y_\xi : \xi < \gamma \rangle \in N$, and we are done.

Claim 4. $\exists$ does not win $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$.

Suppose $\exists$ has a winning strategy τ . Let $\mathfrak{H}$ be defined as above and $\mathfrak{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of $\mathfrak{H}$ of cardinality $\aleph_2$ such that $\alpha = N \cap \omega_3 \in T$. We let $\forall$ play during the first ω moves of $\mathcal{G}_{\omega_1}(\mathfrak{A}, \mathfrak{B})$ a sequence $\langle a_n : n < \omega \rangle$ in $\mathfrak{A}$ such that if α_n is the least α_n with $a_n \in M_{\alpha_n}$, then the sequence $\langle \alpha_n : n < \omega \rangle$ is ascending and $\sup\{\alpha_n : n < \omega\} = \alpha$. Let $\exists$ respond following τ with $\langle b_n : n < \omega \rangle$. As his move number ω player $\forall$ plays some element $a_\omega \in M \setminus M_\alpha$ in $\mathfrak{A}$ and $\exists$ answers according to τ with b_ω .

For all $i \leq \omega$, $R_{(a_0, a_i)}(a_0, a_i)$ holds. Hence $R_{(a_0, a_i)}(b_0, b_i)$ holds. So there is w_i such that $w_i(a_0) = b_0$ and $w_i(a_i) = b_i$. Since $G_{\beta_0}(a_0) = b_0$, by (M5), for each i there is β_i so that $G_{\beta_i}(a_i) = b_i$. We can assume that β_i is chosen to be minimal. Notice that for all i , $\beta_i > \alpha_i$ and for $i < \omega$, $\beta_i \in \mathfrak{N}$. So $\sup\{\beta_i : i < \omega\} = \alpha$.

Also, by the same reasoning as above, for each $i < \omega$, $R_{(a_i, a_\omega)}(b_i, b_\omega)$ holds. Applying (M5), we get that $G_{\beta_\omega}(a_i) = b_i$. Using (M5) again and the minimality of β_i , for all $i < \omega$, $\beta_i \in C_{\beta_\omega}$. Thus α is a limit of elements of C_{β_ω} , contradicting $\alpha \in T$. $\square$

REFERENCES

1. J. Barwise and S. Feferman (eds.), *Model-theoretic languages*, Springer, 1985.
2. S. Ben-David, *On Shelah's compactness of cardinals*, Israel J. Math. **31** (1978), 34–56.
3. M. Burke and M. Magidor, *Shelah's pfc theory and its applications*, Ann. Pure Appl. Logic **50** (1990), 207–254.
4. P. Eklof and A. Mekler, *Almost free modules*, North-Holland, Amsterdam, 1990.
5. F. Galvin, T. Jech, and M. Magidor, *An ideal game*, J. Symbolic Logic **43** (1978), 285–292.
6. M. Gitik and S. Shelah, *Cardinal preserving ideals*, J. Symbolic Logic (to appear).
7. L. Harrington and S. Shelah, *Some exact equiconsistency results in set theory*, Notre Dame J. Formal Logic **26** (1985), 178–188.
8. P. Hill, *New criteria for freeness in abelian groups. II*, Trans. Amer. Math. Soc. **196** (1974), 229–308.
9. W. Hodges and S. Shelah, *Infinite games and reduced products*, Ann. Math. Logic **20** (1981), 77–108.
10. T. Hyttinen, *Games and infinitary languages*, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes **64** (1987), 1–32.
11. T. Jech, *Set theory*, Academic Press, 1978.
12. T. Jech, M. Magidor, W. Mitchell, and K. Prikry, *Precipitous ideals*, J. Symbolic Logic **45** (1980), 1–8.
13. T. Jech and S. Shelah, *Full reflection of stationary sets below $\aleph_\omega$* , J. Symbolic Logic **55** (1990), 822–830.
14. R. Jensen, *The fine structure of the constructible hierarchy*, Ann. Math. Logic **4** (1972), 229–308.
15. M. Nadel and J. Stavi, *$L_{\infty\lambda}$ -equivalence, isomorphism and potential isomorphism*, Trans. Amer. Math. Soc. **236** (1978), 51–74.
16. J.-P. Serre, *Trees*, Springer, 1980.
17. S. Shelah, *Generalized quantifiers and compact logic*, Trans. Amer. Math. Soc. **204** (1975), 342–364.
18. —, *Reflecting stationary sets and successors of singular cardinals*, Arch. Math. Logic **31** (1991), 25–53.
19. —, *Classification theory*, rev. ed., North-Holland, 1990.

DEPARTMENT OF MATHEMATICS, SIMON FRASER UNIVERSITY, BURNABY, BRITISH COLUMBIA,
CANADA

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, ISRAEL

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HELSINKI, HELSINKI, FINLAND

LINKED CITATIONS

- Page 1 of 2 -



You have printed the following article:

The Ehrenfeucht-Fraisse-Game of Length #1

Alan Mekler; Saharon Shelah; Jouko Väänänen

Transactions of the American Mathematical Society, Vol. 339, No. 2. (Oct., 1993), pp. 567-580.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28199310%29339%3A2%3C567%3ATEOL%3E2.0.CO%3B2-8>

This article references the following linked citations. If you are trying to access articles from an off-campus location, you may be required to first logon via your library web site to access JSTOR. Please visit your library's website or contact a librarian to learn about options for remote access to JSTOR.

References

⁵ **An Ideal Game**

F. Galvin; T. Jech; M. Magidor

The Journal of Symbolic Logic, Vol. 43, No. 2. (Jun., 1978), pp. 284-292.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28197806%2943%3A2%3C284%3AAIG%3E2.0.CO%3B2-G>

⁸ **New Criteria for Freeness in Abelian Groups. II**

Paul Hill

Transactions of the American Mathematical Society, Vol. 196. (Sep., 1974), pp. 191-201.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28197409%29196%3C191%3ANCFFIA%3E2.0.CO%3B2-M>

¹² **Precipitous Ideals**

T. Jech; M. Magidor; W. Mitchell; K. Prikry

The Journal of Symbolic Logic, Vol. 45, No. 1. (Mar., 1980), pp. 1-8.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28198003%2945%3A1%3C1%3API%3E2.0.CO%3B2-1>

¹³ **Full Reflection of Stationary Sets Below # #**

Thomas Jech; Saharon Shelah

The Journal of Symbolic Logic, Vol. 55, No. 2. (Jun., 1990), pp. 822-830.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28199006%2955%3A2%3C822%3AFROSSB%3E2.0.CO%3B2-8>

NOTE: The reference numbering from the original has been maintained in this citation list.

LINKED CITATIONS

- Page 2 of 2 -



¹⁵ **L ## -Equivalence, Isomorphism and Potential Isomorphism**

Mark Nadel; Jonathan Stavi

Transactions of the American Mathematical Society, Vol. 236. (Feb., 1978), pp. 51-74.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28197802%29236%3C51%3AIAPI%3E2.0.CO%3B2-O>

¹⁷ **Generalized Quantifiers and Compact Logic**

Saharon Shelah

Transactions of the American Mathematical Society, Vol. 204. (Apr., 1975), pp. 342-364.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28197504%29204%3C342%3AGQACL%3E2.0.CO%3B2-G>