



Automorphism Groups of Homogeneous Structures

Andrés Villaveces - *Universidad Nacional de Colombia - Bogotá*

Arctic Set Theory Workshop 3 - Kilpisjärvi - January 2017

CONTENTS

Reconstructing models

The reconstruction problem

The Small Index Property

SIP beyond first order

Uncountable models, still First Order

SIP (non-elementary)

The setting: strong amalgamation classes

Genericity and Amalgamation Bases

Examples: quasiminimal classes, the Zilber field, j-invariants


A “CLASSICAL ENIGMA”: RECONSTRUCTING FROM SYMMETRY.

At a very classical
extreme, there is the old
enigma:
There is some object M .

A “CLASSICAL ENIGMA”: RECONSTRUCTING FROM SYMMETRY.

At a very classical extreme, there is the old enigma:

There is some object M .
I give you the symmetries of M , i.e. $\text{Aut}(M)$.



A “CLASSICAL ENIGMA”: RECONSTRUCTING FROM SYMMETRY.


At a very classical extreme, there is the old enigma:

There is some object M .

I give you the symmetries of M , i.e.

$\text{Aut}(M)$.

Tell me what is M !



RECONSTRUCTING MODELS?

In Model Theory (and in other parts of Mathematics!), the same naïve enigma has important variants. The main version is usually called “The Reconstruction Problem”:

RECONSTRUCTING MODELS?

In Model Theory (and in other parts of Mathematics!), the same naïve enigma has important variants. The main version is usually called “The Reconstruction Problem”:

- ▶ if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about M ? (In general, not much! by e.g. Ehrenfeucht-Mostowski).

RECONSTRUCTING MODELS?

In Model Theory (and in other parts of Mathematics!), the same naïve enigma has important variants. The main version is usually called “The Reconstruction Problem”:

- ▶ if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about M ? (In general, not much! by e.g. Ehrenfeucht-Mostowski).
- ▶ a more reasonable question: if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about $\text{Th}(M)$?

RECONSTRUCTING MODELS?

In Model Theory (and in other parts of Mathematics!), the same naïve enigma has important variants. The main version is usually called “The Reconstruction Problem”:

- ▶ if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about M ? (In general, not much! by e.g. Ehrenfeucht-Mostowski).
- ▶ a more reasonable question: if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about $\text{Th}(M)$?
- ▶ an even more reasonable question: if for some (FO) structure M we are given $\text{Aut}(M)$, when can we recover **all models biinterpretable with M** ?

RECONSTRUCTING MODELS?

In Model Theory (and in other parts of Mathematics!), the same naïve enigma has important variants. The main version is usually called “The Reconstruction Problem”:

- ▶ if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about M ? (In general, not much! by e.g. Ehrenfeucht-Mostowski).
- ▶ a more reasonable question: if for some (First Order) structure M we are **given** $\text{Aut}(M)$, what can we say about $\text{Th}(M)$?
- ▶ an even more reasonable question: if for some (FO) structure M we are given $\text{Aut}(M)$, when can we recover **all models biinterpretable with M** ?
- ▶ we follow ONE line of reconstruction, different from (but related to) the work of M. Rubin!

WHERE ELSE IN MATHEMATICS?

The “naïve question” is quite important: What information about a model M and $Th(M)$ is contained in the group $Aut(M)$?

WHERE ELSE IN MATHEMATICS?

The “naïve question” is quite important: What information about a model M and $Th(M)$ is contained in the group $Aut(M)$?

What information on a metric structure (M, d, \dots) is contained in the isometry group $Iso(M, d, \dots)$?

WHERE ELSE IN MATHEMATICS?

The “naïve question” is quite important: What information about a model M and $Th(M)$ is contained in the group $Aut(M)$?

What information on a metric structure (M, d, \dots) is contained in the isometry group $Iso(M, d, \dots)$?

- (Anabelian geometry) the [anabelian question](#): recover the isomorphism class of a variety X from its étale fundamental group $\pi_1(X)$. Neukirch, Uchida, for algebraic number fields.

WHERE ELSE IN MATHEMATICS?

The “naïve question” is quite important: What information about a model M and $Th(M)$ is contained in the group $Aut(M)$?

What information on a metric structure (M, d, \dots) is contained in the isometry group $Iso(M, d, \dots)$?

- ▶ (Anabelian geometry) the [anabelian question](#): recover the isomorphism class of a variety X from its étale fundamental group $\pi_1(X)$. Neukirch, Uchida, for algebraic number fields.
- ▶ (Koenigsmann) K and $G_{K(t)/K}$ are biinterpretable for K a perfect field with finite extensions of degree > 2 and prime to $char(K)$.

WHERE ELSE IN MATHEMATICS?

The “naïve question” is quite important: What information about a model M and $Th(M)$ is contained in the group $Aut(M)$?

What information on a metric structure (M, d, \dots) is contained in the isometry group $Iso(M, d, \dots)$?

- ▶ (Anabelian geometry) the [anabelian question](#): recover the isomorphism class of a variety X from its étale fundamental group $\pi_1(X)$. Neukirch, Uchida, for algebraic number fields.
- ▶ (Koenigsmann) K and $G_{K(t)/K}$ are biinterpretable for K a perfect field with finite extensions of degree > 2 and prime to $char(K)$.

These are versions of the same kind of problem - but we will not concentrate on these today. They may however be amenable to model theoretic treatment.

LA RECONSTRUCTION DE STRUCTURES À LA LASCAR

- Every automorphism of M extends uniquely to an automorphism of M^{eq} ; therefore, $Aut(M) \approx Aut(M^{eq})$ canonically.

LA RECONSTRUCTION DE STRUCTURES À LA LASCAR

- ▶ Every automorphism of M extends uniquely to an automorphism of M^{eq} ; therefore, $Aut(M) \approx Aut(M^{eq})$ canonically.
- ▶ Having that $M^{eq} \approx N^{eq}$ implies that M and N are bi-interpretable.

LA RECONSTRUCTION DE STRUCTURES À LA LASCAR

- ▶ Every automorphism of M extends uniquely to an automorphism of M^{eq} ; therefore, $Aut(M) \approx Aut(M^{eq})$ canonically.
- ▶ Having that $M^{eq} \approx N^{eq}$ implies that M and N are bi-interpretable.
- ▶ If M is \aleph_0 -categorical, any open subgroup of $Aut(M)$ is a stabilizer $Aut_\alpha(M)$ for some imaginary α . Also $Aut(M) \curvearrowright \{H \leq Aut(M) \mid H \text{ open}\}$ (conjugation).

LA RECONSTRUCTION DE STRUCTURES À LA LASCAR

- ▶ Every automorphism of M extends uniquely to an automorphism of M^{eq} ; therefore, $Aut(M) \approx Aut(M^{eq})$ canonically.
- ▶ Having that $M^{eq} \approx N^{eq}$ implies that M and N are bi-interpretable.
- ▶ If M is \aleph_0 -categorical, any open subgroup of $Aut(M)$ is a stabilizer $Aut_\alpha(M)$ for some imaginary α . Also $Aut(M) \curvearrowright \{H \leq Aut(M) \mid H \text{ open}\}$ (conjugation).
- ▶ The action $Aut(M) \curvearrowright$ is (almost) \approx to $Aut(M) \curvearrowright M^{eq}$.

LA RECONSTRUCTION DE STRUCTURES À LA LASCAR


- ▶ Every automorphism of M extends uniquely to an automorphism of M^{eq} ; therefore, $Aut(M) \approx Aut(M^{eq})$ canonically.
- ▶ Having that $M^{eq} \approx N^{eq}$ implies that M and N are bi-interpretable.
- ▶ If M is \aleph_0 -categorical, any open subgroup of $Aut(M)$ is a stabilizer $Aut_\alpha(M)$ for some imaginary α . Also $Aut(M) \curvearrowright \{H \leq Aut(M) \mid H \text{ open}\}$ (conjugation).
- ▶ The action $Aut(M) \curvearrowright$ is (almost) \approx to $Aut(M) \curvearrowright M^{eq}$.

So, we have recovered the action of $Aut(M)$ on M^{eq} from the topology of $Aut(M)$... so, if M, N are countable \aleph_0 -categorical structures, TFAE:

- ▶ There is a bicontinuous isomorphism from $Aut(M)$ onto $Aut(N)$
- ▶ M and N are bi-interpretable.


SOME (NON-)EXAMPLES - WHY IS SATURATION NEEDED?

Let M_1 be the countable
saturated model of P_i ($i < \omega$)
disjoint infinite predicates and




SOME (NON-)EXAMPLES - WHY IS SATURATION NEEDED?

Let M_1 be the countable saturated model of P_i ($i < \omega$) disjoint infinite predicates and




let M_2 be a countable model of an equivalence relation with infinitely many infinite classes, with exactly one constant c_i in each class but one




SOME (NON-)EXAMPLES - WHY IS SATURATION NEEDED?

Let M_1 be the countable saturated model of P_i ($i < \omega$) disjoint infinite predicates and




let M_2 be a countable model of an equivalence relation with infinitely many infinite classes, with exactly one constant c_i in each class but one




yet $Aut(M_1) \approx Aut(M_2)$

SIP - THE LINK BETWEEN ALGEBRA AND TOPOLOGY



SIP - THE LINK BETWEEN ALGEBRA AND TOPOLOGY



Now, to the main property of the group $Aut(M)$ that enables us to capture its topology...

THE SMALL INDEX PROPERTY (COUNTABLE VERSION)

Definition (Small Index Property - SIP)

Let M be a countable structure. M has the small index property if for any subgroup H of $\text{Aut}(M)$ of index less than 2^{\aleph_0} , there exists a finite set $A \subset M$ such that $\text{Aut}_A(M) \subset H$.

BASIC FACTS ON COUNTABLE SIP

SIP allows us to recover the topological structure of $\text{Aut}(M)$ from its pure group structure:

Open neighborhoods of 1 in pointwise convergence topology =

Subgroups containing pointwise stabilizers $\text{Aut}_A(M)$ for some finite A .

BASIC FACTS ON COUNTABLE SIP

SIP allows us to recover the topological structure of $\text{Aut}(M)$ from its pure group structure:

Open neighborhoods of 1 in pointwise convergence topology =

Subgroups containing pointwise stabilizers $\text{Aut}_A(M)$ for some finite A .

- ▶ SIP holds for random graph, infinite set, DLO, vector spaces over finite fields, generic relational structures, \aleph_0 -categorical \aleph_0 -stable structures, etc.
- ▶ It fails e.g. for $M \models \text{ACF}_0$ with ∞ transc. degree.

GALOIS GROUP (OF A THEORY)


The Galois group of a model M ,

$$\text{Gal}(M) := \text{Aut}(M) / \text{Aut}_f(M),$$

is invariant across saturated models of a theory.

Possible failures of SIP are encoded in this quotient.

TO THE UNCOUNTABLE / THE NON-ELEMENTARY



SIP FOR UNCOUNTABLE STRUCTURES

We now switch focus to the uncountable, first order, case.

Fix $\lambda = \lambda^{<\lambda}$ an uncountable cardinal, and fix M a saturated model of cardinality λ .

SIP FOR UNCOUNTABLE STRUCTURES

We now switch focus to the uncountable, first order, case.

Fix $\lambda = \lambda^{<\lambda}$ an uncountable cardinal, and fix M a saturated model of cardinality λ .

We now use the topology \mathcal{T}^λ on $\text{Aut}(M)$, whose basic open sets around 1_M are stabilizers of subsets of size $< \lambda$ - as before $\text{Aut}_A(M)$ but now $A \subset M$ with $|A| < \lambda$.

SIP FOR UNCOUNTABLE STRUCTURES

We now switch focus to the uncountable, first order, case.

Fix $\lambda = \lambda^{<\lambda}$ an uncountable cardinal, and fix M a saturated model of cardinality λ .

We now use the topology \mathcal{T}^λ on $\text{Aut}(M)$, whose basic open sets around 1_M are stabilizers of subsets of size $< \lambda$ - as before $\text{Aut}_A(M)$ but now $A \subset M$ with $|A| < \lambda$.

$\text{Aut}(M)$ with this topology is of course no longer a Polish space. The techniques from Descriptive Set Theory that have been used for the countable case need to be replaced (Friedman, Hyttinen and Kulikov have a start of Descriptive Set Theory for some uncountable cardinalities, however).

LASCAR-SHELAH'S THEOREM

Theorem (Lascar-Shelah: Uncountable saturated models have the SIP)

Let M be saturated, of cardinality $\lambda = \lambda^{<\lambda}$ and let G be a subgroup of $\text{Aut}(M)$ such that $[\text{Aut}(M) : G] < 2^\lambda$. Then there exists $A \subset M$ with $|A| < \lambda$ such that $\text{Aut}_A(M) \subset G$.

LASCAR-SHELAH'S THEOREM

Theorem (Lascar-Shelah: Uncountable saturated models have the SIP)

Let M be saturated, of cardinality $\lambda = \lambda^{<\lambda}$ and let G be a subgroup of $\text{Aut}(M)$ such that $[\text{Aut}(M) : G] < 2^\lambda$. Then there exists $A \subset M$ with $|A| < \lambda$ such that $\text{Aut}_A(M) \subset G$.

The proof consists of building directly (assuming that G does not contain any open set $\text{Aut}_A(M)$ around the identity) a **binary tree** of height λ of automorphisms of M in such a way that every two of them are not conjugate. This is enough but requires two crucial notions: **generic** and **existentially closed (sequences of automorphisms)**. These are obtained by assuming that G is not open.

BEYOND FIRST ORDER

Although results on the reconstruction problem, so far have been stated and proved for saturated models in first order theories, the scope of the matter can go far beyond:

- Abstract Elementary Classes with well-behaved closure notions, and the particular case:

BEYOND FIRST ORDER

Although results on the reconstruction problem, so far have been stated and proved for saturated models in first order theories, the scope of the matter can go far beyond:

- ▶ Abstract Elementary Classes with well-behaved closure notions, and the particular case:
- ▶ Quasiminimal (qm excellent) Classes.

THE SETTING: STRONG AMALGAMATION CLASSES

A setting for homogeneity: let $(\mathcal{K}, \prec_{\mathcal{K}})$ be an AEC, with $LS(\mathcal{K}) \leq \lambda$, $|M| = \kappa > \lambda$, $\kappa^{<\kappa} = \kappa$.

Let $\mathcal{K}^{<}(M) := \{N : N \preceq_K M, |N| < \kappa\}$ and fix $M \in \mathcal{K}$ homogeneous.

THE SETTING: STRONG AMALGAMATION CLASSES

A setting for homogeneity: let $(\mathcal{K}, \prec_{\mathcal{K}})$ be an AEC, with $LS(\mathcal{K}) \leq \lambda$, $|M| = \kappa > \lambda$, $\kappa^{<\kappa} = \kappa$.

Let $\mathcal{K}^{<}(M) := \{N : N \preceq_K M, |N| < \kappa\}$ and fix $M \in \mathcal{K}$ homogeneous.

The topology τ^{cl} : base of open neighborhoods given by sets of the form $Aut_X(M)$ where $X \in \mathcal{C}$, where

$\mathcal{C} := \{cl^M(A) : A \subseteq M \text{ such that } |A| < \kappa\}$ and the “closure operator” is $cl^M(A) := \bigcap_{A \subset N \prec_{\mathcal{K}} M} A$.

This class of cl^M -closed sets has enough structure for the proof of SIP.

THE MAIN RESULT: SIP FOR HOMOGENEOUS AEC.

Theorem (SIP for $(\text{Aut}(M), \mathcal{T}^{cl})$ - Ghadernezhad, V.)

Let M be a homogeneous model in an AEC $(\mathcal{K}, \prec_{\mathcal{K}})$, with $|M| = \lambda = \lambda^{<\lambda} > LS(\mathcal{K})$, such that $\mathcal{K}^{<\lambda}$ is a strong amalgamation class. Let $G \leq \text{Aut}(M)$ with $[\text{Aut}(M) : G] \leq \lambda$ (this is, G has small index in $\text{Aut}(M)$). Then there exists $X \in \mathcal{C}$ such that $\text{Aut}_X(M) \leq G$ (i.e., G is open in $(\text{Aut}(M), \mathcal{T}^{cl})$).

GETTING MANY NON-CONJUGATES

Proof (rough sketch): suppose G has small index in $\text{Aut}(M)$ but is not open (does not contain any basic $\text{Aut}_X(M)$ for $X \in \mathcal{C}$).

We have enough tools (**generic** sequences and **strong amalgamation bases**) to build a Lascar-Shelah tree to reach a contradiction (2^λ many branches giving automorphisms of M g_σ for $\sigma \in 2^\lambda$ such that if $\sigma \neq \tau \in 2^\lambda$ then $g_\sigma^{-1} \circ g_\tau \notin G$).

Of course, the possibility of getting these 2^λ -many automorphisms requires using the non-openness of G to get the construction going.

LASCAR-SHELAH TREE FOR OUR SITUATION

A λ -Lascar-Shelah tree for M and $G \leq \text{Aut}(M)$ is a binary tree of height λ with, for each $s \in 2^{<\lambda}$, a model $M_s \in \mathcal{K}^{<}(M)$, $g_s \in \text{Aut}(M_s)$, $h_s, k_s \in \text{Aut}_{M_s}(M)$ such that

- ▶ $h_{s,0} \in G$ and $h_{s,1} \notin G$ for all $s \in \mathcal{S}$;
- ▶ $k_{s,0} = k_{s,1}$ for all $s \in \mathcal{S}$;
- ▶ for $s \in \mathcal{S}$ and all $t \in \mathcal{S}$ such that $t \leq s : h_t \restriction [M_s] = M_s$ (i.e. $h_t \in \text{Aut}_{\{M_s\}}(M)$) and ...;
- ▶ for $s \in \mathcal{S}$ and all $t \in \mathcal{S}$ such that $t \leq s : g_s \cdot (h_t \restriction M_s) \cdot g_s^{-1} = k_t \restriction M_s$;
- ▶ for $s \in \mathcal{S}$ and $\beta < \text{length}(s)$: $a_s \in M_s$;
- ▶ for all s , the families $\{h_t : t \leq s, t \in \mathcal{S}\}$ and $\{k_t : t \leq s, t \in \mathcal{S}\}$ are elements of \mathcal{F} (i.e. they are generic).

GENERIC SEQUENCES AND STRONG AMALGAMATION BASES

The main technical tools in the construction of a LS tree are

- Guaranteeing generic sequences of automorphisms

$(g \in \text{Aut}(M))$ is generic if

$\forall N \in \mathcal{K}^{<}(M)$ such that $g \upharpoonright N \in \text{Aut}(N)$

$\forall N_1 \succ_{\mathcal{K}} N, N_1 \in \mathcal{K}^{<}(M)$

$\forall h \supset g \upharpoonright N, h \in \text{Aut}(N_1)$

$\exists \alpha \in \text{Aut}_N(M)$ such that $g \supset \alpha \circ h \circ \alpha^{-1}$,

- showing they are unique up to conjugation,
- getting a generic sequence $\mathcal{F} = (g_i : i \in I)$ such that
 1. the set $\{i \in I : g_i \upharpoonright M_0 = h \text{ and } g_i \notin G\}$ has cardinality κ for all $M_0 \in \mathcal{K}^{<}(M)$ and $h \in \text{Aut}(M_0)$;
 2. the set $\{i \in I : g_i \in G\}$ has cardinality κ .

QUASIMINIMAL PREGEOMETRY CLASSES

In a language L , a quasiminimal pregeometry class \mathcal{Q} is a class of pairs $\langle H, \text{cl}_H \rangle$ where H is an L -structure, cl_H is a pregeometry operator on H such that the following conditions hold:

1. Closed under isomorphisms,
2. For each $\langle H, \text{cl}_H \rangle \in \mathcal{Q}$, the closure of any finite set is countable.
3. If $\langle H, \text{cl}_H \rangle \in \mathcal{Q}$ and $X \subseteq H$, then $\langle \text{cl}_H(X), \text{cl}_H \upharpoonright \text{cl}_H(X) \rangle \in \mathcal{Q}$.
4. If $\langle H, \text{cl}_H \rangle, \langle H', \text{cl}_{H'} \rangle \in \mathcal{Q}$, $X \subseteq H$, $y \in H$ and $f : H \rightarrow H'$ is a partial embedding defined on $X \cup \{y\}$, then $y \in \text{cl}_H(X)$ if and only if $f(y) \in \text{cl}_{H'}(f(X))$.
5. Homogeneity over countable models.

QUASIMINIMAL PREGEOMETRY CLASSES

In a language L , a quasiminimal pregeometry class \mathcal{Q} is a class of pairs $\langle H, \text{cl}_H \rangle$ where H is an L -structure, cl_H is a pregeometry operator on H such that the following conditions hold:

1. Closed under isomorphisms,
2. For each $\langle H, \text{cl}_H \rangle \in \mathcal{Q}$, the closure of any finite set is countable.
3. If $\langle H, \text{cl}_H \rangle \in \mathcal{Q}$ and $X \subseteq H$, then $\langle \text{cl}_H(X), \text{cl}_H \upharpoonright \text{cl}_H(X) \rangle \in \mathcal{Q}$.
4. If $\langle H, \text{cl}_H \rangle, \langle H', \text{cl}_{H'} \rangle \in \mathcal{Q}$, $X \subseteq H$, $y \in H$ and $f : H \rightarrow H'$ is a partial embedding defined on $X \cup \{y\}$, then $y \in \text{cl}_H(X)$ if and only if $f(y) \in \text{cl}_{H'}(f(X))$.
5. Homogeneity over countable models.

These can all be generated by ONE canonical structure.

VERY RECENT UPDATES

- ▶ In November 2016 - about a week ago, Sébastien Vasey has posted a paper on the ArXiv proving that quasiminimal pregeometries do not require the exchange axiom of pregeometries. This makes it in principle easier to prove that classes are quasiminimal!
- ▶ Vasey has also suggested that our theorem applies to wider classes (excellent classes, and even wider: certain “non-forking frames”). This is work in progress now.

EXAMPLE: QUASIMINIMAL CLASSES, “ZILBER FIELD”

- \mathcal{Q} quasiminimal pregeometry class. $M \in \mathcal{Q}$ of size \aleph_1 , $\mathcal{C} = \{cl(A) \mid A \subset M, A \text{ small}\}$ then \mathcal{C} has the free aut-independence amalgamation property. (Based on Haykazyan's paper on qm classes.)

EXAMPLE: QUASIMINIMAL CLASSES, “ZILBER FIELD”

- ▶ \mathcal{Q} quasiminimal pregeometry class. $M \in \mathcal{Q}$ of size \aleph_1 , $\mathcal{C} = \{cl(A) \mid A \subset M, A \text{ small}\}$ then \mathcal{C} has the free aut-independence amalgamation property. (Based on Haykazyan's paper on qm classes.)
- ▶ \mathcal{Q} qm pregeom. class \rightarrow for every model M of \mathcal{Q} , $Aut(M)$ has SIP,

EXAMPLE: QUASIMINIMAL CLASSES, “ZILBER FIELD”

- ▶ \mathcal{Q} quasiminimal pregeometry class. $M \in \mathcal{Q}$ of size \aleph_1 , $\mathcal{C} = \{cl(A) \mid A \subset M, A \text{ small}\}$ then \mathcal{C} has the free aut-independence amalgamation property. (Based on Haykazyan’s paper on qm classes.)
- ▶ \mathcal{Q} qm pregeom. class \rightarrow for every model M of \mathcal{Q} , $Aut(M)$ has SIP,
- ▶ The “Zilber field” has SIP.

EXAMPLE: QUASIMINIMAL CLASSES, “ZILBER FIELD”

- ▶ \mathcal{Q} quasiminimal pregeometry class. $M \in \mathcal{Q}$ of size \aleph_1 , $\mathcal{C} = \{cl(A) \mid A \subset M, A \text{ small}\}$ then \mathcal{C} has the free aut-independence amalgamation property. (Based on Haykazyan’s paper on qm classes.)
- ▶ \mathcal{Q} qm pregeom. class \rightarrow for every model M of \mathcal{Q} , $Aut(M)$ has SIP,
- ▶ The “Zilber field” has SIP.
- ▶ The j -invariant has the SIP.

REFERENCES



Friedman, Sy; Hyttinen, Tapani and Kulikov, Vadim Descriptive Set Theory on Uncountable...,



Lascar, Daniel. Automorphism Groups of Saturated Structures, ICM 2002, Vol. III - 1-3.



Lascar, Daniel. Les automorphismes d'un ensemble fortement minimal, JSL, vol. 57, n. 1. March 1992.



Lascar, Daniel and Shelah, Saharon. Uncountable Saturated Structures have the Small Index Property, Bull. London Math. Soc. 25 (1993) 125-13.1



Ghadernezhad, Zaniar and Villaveces, Andrés. The Small Index Property for Homogeneous Abstract Elementary Classes, submitted (2016) to IPM 2015 Proceedings.



Thanks, thanks, to you all (and especially to J and J)