

Open dihypergraphs on generalized Baire spaces

Philipp Schlicht, University of Bristol

Arctic Set Theory Workshop 5, Kilpisjärvi, 17 February 2022

Based on a joint project with Dorottya Sziráki, Budapest.

- Philipp Schlicht, Dorottya Sziráki:
The open dihypergraph dichotomy for generalized Baire spaces,
70 pages, in preparation

Motivation: Ramsey theory

It is natural to wonder whether Ramsey's theorem for n -tuples of natural numbers can be extended to the set of real numbers.

- Sierpinski's counterexample 1937: a partition of pairs of reals in two pieces with no uncountable homogeneous set
- Galvin 1968: Ramsey's theorem for open graphs on the reals
- Blass 1981: A generalization to Borel n -hypergraphs on the reals

Motivation: Ramsey theory

Galvin's theorem can be strengthened.

- Todorćević's **open graph axiom** 1989:
- Feng's **open graph dichotomy** for analytic sets 1993:

Feng's theorem implies one of the most basic descriptive set theoretic dichotomies: the perfect set property.

The graph-theoretic approach

In the last few years, graph dichotomies provided new proofs of old and new theorems in descriptive set theory.

Kechris, Solecki, Todorcevic and Miller proved results for analytic graphs (variants of the G_0 -dichotomy) that imply:

- Suslin's perfect set property of analytic sets
- Lusin and Novikov's uniformization of Borel sets with countable sections
- Feng's open graph dichotomy
- Silver's theorem on coanalytic equivalence relations

The graph-theoretic approach

Carroy, Miller and Soukup 2020 found an **infinite dimensional** version of Feng's open graph dichotomy.

Note the following restrictions:

- Farah, Todorcevic 1995: The open graph dichotomy fails for closed graphs.
- Farah, Todorcevic 1995, He 2005: The open 3-hypergraph dichotomy fails.

One thus has to consider **directed** hypergraphs.

A **κ -dihypergraph** on X is a set of nonconstant sequences in ${}^\kappa X$.

The open graph dichotomy

A **graph** G is a symmetric relation with no loops.

A graph G on a space X is an **open graph** if it is an open subset of $X \times X$ without the diagonal.

Definition (Feng 1993)

$\text{OGD}_\omega(X)$ states that for any open graph G on X , either

1. G has an **ω -coloring** or
2. G has a **perfect complete subgraph**.

G has an **ω -coloring** if and only if X is the union of countably many G -independent sets.

A high dimensional dichotomy

We fix the **box topology** on ${}^\omega X$ with basic open sets $\prod_{i < \omega} U_i$, where each U_i is open in X .

Definition (Carroy, Miller, Soukup 2020)

$\text{ODD}_\omega^\omega(X)$ states that for any box-open ω -**dihypergraph** H on X , either

1. H has a ω -**coloring** or
2. there is a **continuous homomorphism** $f: {}^\omega \omega \rightarrow X$ from $\mathbb{H}_{\omega\omega}$ to H .

$$\mathbb{H}_{\omega\omega} = \{ \vec{x} \in {}^\omega({}^\omega \omega) \mid \exists t \in {}^{<\omega} \omega \ \forall n \in \omega \ t \frown \langle n \rangle \subseteq x_n \}$$

$\text{ODD}_\omega^\omega(X, H)$ states that this holds for H .

Theorem (CMS)

$\text{ODD}_\omega^\omega(X)$ holds for all analytic subsets X of ${}^\omega\omega$.

It holds for all subsets, assuming AD.

They prove a number of applications:

1. The **Hurewicz dichotomy** for X : either
 - X is contained in a K_σ set, or
 - X contains a closed subset homeomorphic to ${}^\omega\omega$.
2. The **Jayne-Rogers theorem** on piecewise continuous functions with closed pieces on X .
3. A theorem of Lecomte and Zeleny on Δ_2^0 -measurable ω -colorings on X .

Applications

For a metric space X , let H_X denote the ω -dihypergraph on X of all **injective** sequences in X with **no convergent subsequence**.

Proposition (CMS)

1. H_X is box-open.
2. There is an ω -coloring of $H_X \upharpoonright Y$ iff Y is contained in a K_σ set.
3. A continuous function ${}^\omega\omega \rightarrow X$ is a **homomorphism** from $\mathbb{H}_{\omega\omega}$ to H_X iff it is an **injective closed** map.

Proof sketch.

For 2., note that a subset Y of X is H_X -independent iff its closure is **compact**. □

Generalized Baire spaces

κ always denotes an uncountable cardinal with $\kappa^{<\kappa} = \kappa$.

Definitions are analogous:

- The κ -Baire space ${}^\kappa\kappa$ is the set of functions $x : \kappa \rightarrow \kappa$ with the bounded topology. The basic open sets are

$$N_t = \{x \in {}^\kappa\kappa \mid t \subseteq x\}$$

for all $t \in {}^{<\kappa}\kappa$.

- The κ -Cantor space ${}^\kappa 2$ has subspace topology.
- κ -Borel sets are generated from open sets by closing under unions and intersections of size κ and negations.
- κ -analytic sets are continuous images of closed sets.

Relative to an inaccessible cardinal:

Theorem (Lücke, Motto Ros, S. 2016)

*The **Hurewicz dichotomy** for all κ -analytic subsets of ${}^\kappa\kappa$ is consistent.*

Theorem (S. 2017)

*The **perfect set property (PSP)** for all **definable subsets** of ${}^\kappa\kappa$ is consistent.*

By **definable** we mean definable from a sequence in ${}^\kappa\text{Ord}$.

From OGD to ODD

Theorem (Sziraki 2018)

The *open graph dichotomy (OGD)* for all κ -analytic subsets of ${}^\kappa\kappa$ is consistent.

Definition

$\text{ODD}_\kappa^\kappa(X)$ states that for any box-open κ -dihypergraph H on X , either

1. H has a κ -coloring or
2. there is a *continuous homomorphism* $f: {}^\kappa\kappa \rightarrow X$ from \mathbb{H}_κ to H .

$$\mathbb{H}_\kappa = \{ \vec{x} \in {}^\kappa({}^\kappa\kappa) \mid \exists t \in <{}^\kappa\kappa \ \forall i \in \kappa \ t \frown \langle i \rangle \subseteq x_i \}$$

$\text{ODD}_\kappa^\kappa(X, H)$ states that this holds for H .

ODD_κ^α denotes the version for α -dihypergraphs.

Theorem (Sziraki, S. 2021)

Suppose that V is a $\text{Col}(\kappa, <\lambda)$ -generic extension. Then

$\text{ODD}_\omega^\omega(X, H)$ holds for all *definable* subsets X of ${}^\kappa\kappa$ and:

1. *all definable* box-open κ -dihypergraphs H on X , if λ is *inaccessible* in the ground model.
2. *arbitrary* box-open κ -dihypergraphs H on X , if λ is *Mahlo* in the ground model.

- All applications of CMS in the countable case are consistent relative to an inaccessible or Mahlo cardinal. They do not need AD.
- The Hurewicz dichotomy: X contains a closed homeomorphic copy of ${}^\kappa\kappa$ or X is contained in a union of κ many κ -compact sets.

Example

$\text{ODD}_{\kappa}^2(X)$ implies the open graph dichotomy $\text{OGD}_{\kappa}(X)$.

To see this, take $x \neq y$ in ${}^{\kappa}2$. Let $i < \kappa$ be least with $x(i) \neq y(i)$.

$$\langle x, y \rangle \in \mathbb{H}_{\kappa_2} \iff x(i) = 0 \wedge y(i) = 1.$$

The complete graph \mathbb{K}_{κ_2} on ${}^{\kappa}2$ is the smallest (symmetric) graph containing \mathbb{H}_{κ_2} .

Thus a continuous **homomorphism** $f: {}^{\kappa}2 \rightarrow X$ from \mathbb{H}_{κ_2} to a graph G is also a homomorphism from \mathbb{K}_{κ_2} to G .

Note that f is injective. So G has a perfect complete subgraph.

Step 1: Reflection

Notation: Let G be $\text{Col}(\kappa, <\lambda)$ -generic, where $\lambda > \kappa$ is inaccessible.

For each $\alpha < \lambda$, let $G_\alpha = G \cap \text{Col}(\kappa, <\alpha)$. Write

$$X_{\varphi,a} = \{x \in {}^\kappa\kappa : \varphi(x, a)\}$$

Lemma

Suppose $X \subseteq {}^\kappa\kappa$. If X is definable in $V[G]$ or λ is Mahlo in V , then

$$X \cap V[G_\nu] \in V[G_\nu]$$

for stationarily many $\nu < \lambda$.

Proof sketch.

If X is definable in $V[G]$, the claim holds for a tail of $\nu < \kappa$, since the tail forcings are homogeneous.

Now suppose that λ is Mahlo in V .

Step 1: Reflection

Let \dot{X} be a name for X . Define $f: \lambda \rightarrow \lambda$ as follows.

For $\alpha < \lambda$ and a nice $\text{Col}(\kappa, <\alpha)$ -name $\dot{x} \in V$ for a subset of $\kappa \times \kappa$, let $A_{\dot{x}}$ be a maximal antichain in $\text{Col}(\kappa, <\lambda)$ deciding $\dot{x} \in \dot{X}$.

Since $\text{Col}(\kappa, <\lambda)$ has the λ -c.c., let $f(\alpha) < \lambda$ be such that $A_{\dot{x}} \subseteq \text{Col}(\kappa, <f(\alpha))$ for all such nice names \dot{x} .

The set S of inaccessible closure points of f is stationary, since λ is Mahlo.

Claim

$X \cap V[G_\nu] \in V[G_\nu]$ for all $\nu \in S$.

Let

$$F_\nu(\dot{x}^{G_\nu}) = \begin{cases} 1 & \text{if } p \Vdash_{\text{Col}(\kappa, <\lambda)}^V \dot{x} \in \dot{X} \text{ for some } p \in G_\nu, \\ 0 & \text{if } p \Vdash_{\text{Col}(\kappa, <\lambda)}^V \dot{x} \notin \dot{X} \text{ for some } p \in G_\nu. \end{cases}$$

F_ν is the characteristic function of $X \cap V[G_\nu]$, since $G_\nu \subseteq G$.

Step 2: Independent trees

In $V[G]$, suppose $a \in {}^\kappa \text{Ord}$. Write

$$X_{\varphi,a} = \{x \in {}^\kappa \kappa \mid \varphi(x, a)\}.$$

$$\mathcal{T}^{\text{ind}} = \{T \subseteq {}^{<\kappa} \kappa \mid T \text{ is a tree, } [T] \text{ is } R\text{-independent}\}.$$

Then $\mathcal{T}^{\text{ind}} \cap V[G_\nu] \in V[G_\nu]$ for some $\nu < \lambda$ with $a \in V[G_\nu]$ by the previous step. We can assume $V[G_\nu] = V$.

If R has no κ -coloring, then for some $\gamma < \lambda$:

$$(X_{\varphi,a} \setminus \bigcup \{[T] \mid T \in \mathcal{T}_V^{\text{ind}}\}) \cap V[G_\gamma] \neq \emptyset.$$

Step 2: Independent trees

In V , let \dot{x} be a $\text{Col}(\kappa, <\gamma)$ -name for an element of $X_{\varphi,a}$ such that $1_{\text{Col}(\kappa, <\gamma)} \Vdash \dot{x} \notin [T]$ for all $T \in \mathcal{T}_V^{\text{ind}}$. For any $p \in \text{Col}(\kappa, <\gamma)$, let

$$T^{\dot{x},p} = \{t \in {}^{<\kappa}\kappa \mid \exists q \leq p \ q \Vdash t \subseteq \dot{x}\}$$

denote the *tree of possible values* for \dot{x} below p .

Lemma

1. $1_{\text{Col}(\kappa, <\gamma)} \Vdash \text{"}\dot{x} \in X_{\varphi,a} \text{ in every further } \text{Col}(\kappa, <\lambda)\text{-gen. extension.}"$
2. $T^{\dot{x},p} \notin \mathcal{T}_V^{\text{ind}}$ for all $p \in \text{Col}(\kappa, <\gamma)$.

Proof of 2. $p \Vdash \dot{x} \in [T^{\dot{x},p}]$. □

We now assume \dot{x} is an $\text{Add}(\kappa, 1)$ -name.

Step 3: Construction of a forcing \mathbb{Q}

The forcing will construct the required homomorphism. The point is to **avoid** subsets of κ with bad **quotients**.

We construct a forcing \mathbb{Q} such that:

1. \mathbb{Q} is **equivalent** to $\text{Add}(\kappa, 1)$.
2. Suppose that $V[H]$ is any \mathbb{Q} -generic extension of V . \mathbb{Q} adds a map $g : (\kappa_\kappa)^{V[H]} \rightarrow (\kappa_\kappa)^{V[H]}$ such that for each $y \in (\kappa_\kappa)^{V[H]}$,
 - $g(y)$ is **$\text{Add}(\kappa, 1)$** -generic over V ,
 - $V[H]$ is a **$\text{Add}(\kappa, 1)$** -generic extension of $V[g(y)]$, and
 - $\dot{x}^{g(y)} \in X_{\varphi, a}$.

$f : \kappa_\kappa \rightarrow X, f(y) = \dot{x}^{g(y)}$ is **continuous**.

3. f is a **homomorphism** from $\mathbb{H}_{\kappa_\kappa}$ to R .

The main work is to prove properties of \mathbb{Q} .

Future directions

Inaccessibles are necessary. Are **Mahlo cardinals** necessary for results? (They are for the proofs.) This would separate the variant for **arbitrary** dihypergraphs from the **definable** variant.

Do other large cardinals play a role for the structure of definable subsets of generalized Baire spaces?

Regarding more complex graphs, one can ask: Is a version of the G_0 -dichotomy for ${}^\kappa\kappa$ consistent? Is the Lusin-Novikov theorem for ${}^\kappa\kappa$ consistent?

Future directions

Can the above dichotomies be separated for different dimensions? Is it consistent that $\text{OGD}_\kappa(X)$ holds, but $\text{ODD}_\kappa^\kappa(X)$ fails? Note that all models above are Levy collapses.

To separate the version with injective homomorphisms for ω_1 , one might consider Jensen's and Jonsbraten's model for the Suslin problem with CH.

We are currently studying several variants of the Hurewicz dichotomy, and other applications of Carroy, Soukup and Miller.


Consider the following versions of $\text{ODD}_{\kappa}^{\kappa}(X, H)$ with the condition on the homomorphism is strengthened.

- $\text{ODDC}_{\kappa}^{\kappa}(X, H)$: homeomorphism onto a closed image
- $\text{ODDH}_{\kappa}^{\kappa}(X, H)$: homeomorphism onto its image
- $\text{ODDI}_{\kappa}^{\kappa}(X, H)$: injective

Variants



The $\kappa = \omega$
case



$\kappa = \omega_1$ or κ is weakly inaccessible but not inaccessible



$\kappa \geq \omega_2$ is a successor cardinal or is inaccessible

The implications from left to right hold by definition.

$A \longleftrightarrow B$: A and B are equivalent for all X, H .

$A \implies B$: A implies B for all X, H and the implication is strict, i.e., there exist X, H such that the reverse implication fails.

solid arrow: provable for all κ with $\kappa^{<\kappa} = \kappa$.

dashed arrow: consistent and follows from the assumption in the superscript.

dotted arrow: its consistency is an open question.

Definition

Let \mathbb{D}_κ denote the κ -dimensional dihypergraph on ${}^\kappa\kappa$ consisting of all non-constant sequences $\langle x_\alpha : \alpha < \kappa \rangle$ which are **dense** in some **basic open** subset of ${}^\kappa\kappa$.¹

Lemma

\mathbb{D}_κ separates $\text{ODDH}_\kappa^\kappa(X, H)$ from $\text{ODD}_\kappa^\kappa(X, H)$.

Lemma

There exists a box-open κ -dimensional dihypergraph H on ${}^\kappa\kappa$ that separates $\text{ODDC}_\kappa^\kappa(X, H)$ from $\text{ODDH}_\kappa^\kappa(X, H)$.

¹i.e., $\{x_\alpha : \alpha < \kappa\} \cap N_t$ is a dense subset of N_t for some $t \in {}^{<\kappa}\kappa$.