

The Calderón Problem for Conormal Potentials I: Global Uniqueness and Reconstruction

ALLAN GREENLEAF

University of Rochester

MATTI LASSAS

University of Helsinki

AND

GUNTHER UHLMANN

University of Washington

1 Introduction

The goal of this paper is to establish global uniqueness and obtain reconstruction, in dimensions $n \geq 3$, for the Calderón problem in the class of potentials conormal to a smooth submanifold H in \mathbb{R}^n . In the case of hypersurfaces, the potentials considered here may have any singularity weaker than that of the delta function δ_H on the hypersurface H ; in general, these potentials correspond to conductivities that are in $C^{1+\epsilon}$ and thus fail to be covered by previously known results.

Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain, $H \subset \Omega$ a smooth submanifold of codimension k , and $q \in I^\mu(H)$ a real conormal distribution of order μ with $\mu < 1 - k$. Thus, if $H = \{x : F_j(x) = 0, 1 \leq j \leq k\}$ is a local representation of H by means of defining functions with $\{\nabla F_j : 1 \leq j \leq k\}$ linearly independent on H , then locally $q(x)$ has the Fourier integral representation

$$(1.1) \quad q(x) = \int_{\mathbb{R}^k} e^{i \sum_j F_j(x) \cdot \theta_j} a(x, \theta) d\theta, \quad a \in S_{1,0}^\mu,$$

where $S_{1,0}^\mu$ denotes the standard class of symbols of order μ and type $(1, 0)$ on $\mathbb{R}^n \times (\mathbb{R}^k \setminus 0)$. (Here we use the order convention of [12] rather than [14].) A general element $q \in I^\mu(H)$ is a locally finite sum of such expressions. We assume throughout that $\text{supp}(q)$ is compact in Ω . If $-k < \mu < 0$, then q satisfies $|q(x)| \leq C \cdot \text{dist}(x, H)^{-k-\mu}$, so that $q \in L^{\frac{k}{k+\mu}-\epsilon}(\Omega) \forall \epsilon > 0$, and no better in general; in particular, a general element of $I^\mu(H)$ is unbounded. For comparison, surface measure $\delta_H \in I^0(H)$ and, in the hypersurface case, a Heaviside discontinuity across H belongs to $I^{-1}(H)$.

Rather than working with the Dirichlet-to-Neumann map Λ_q , we state our main results in terms of the Cauchy data \mathcal{CD}_q of sufficiently regular solutions of the

Schrödinger equation

$$(1.2) \quad (\Delta + q(x))u(x) = 0 \quad \text{on } \Omega.$$

This is more flexible, since \mathcal{CD}_q can be defined for potentials q for which Λ_q is either not defined (for example, if $\lambda = 0$ is a Dirichlet eigenvalue) or not known to be defined (due to the low regularity of $q(x)$); it is perhaps more natural as well.

It will be convenient to write $\mu = \nu - k$. Assume that $\nu_0(k) < \nu < 1$ where

$$\nu_0(k) \stackrel{\text{def}}{=} \max\left(\frac{2}{3}, 1 - \frac{k}{4}\right).$$

Fix p and r satisfying

$$(1.3) \quad 2 \leq r < \frac{k}{2(1-\nu)} \stackrel{\text{def}}{=} r_0(k, \nu) < p_0(k, \nu) \stackrel{\text{def}}{=} \frac{2k}{k-\nu} < p < \infty.$$

(If $\nu \leq \nu_0(k)$, just pick p and r for some $\nu' > \nu_0(k)$.) Fixing a smooth function $\psi \in C^\infty(\mathbb{R}^n)$, $\psi \equiv 1$, near $\partial\Omega$ and $\text{int}\{\psi = 0\} \cap \Omega \neq \emptyset$, define the norm

$$(1.4) \quad \|f\|_{X^{p,r}} = \|f\|_{L^p(\Omega)} + \|\Delta f\|_{L^{p'}(\Omega)} + \|\psi f\|_{W^{2,r}(\Omega)},$$

where p' is the dual exponent to p and $W^{2,r}$ is the standard Sobolev space of $f \in \mathcal{D}'(\Omega)$ having two derivatives in $L^r(\Omega)$. Set

$$(1.5) \quad X^{p,r}(\Omega) = \{f \in \mathcal{D}'(\Omega) : \|f\|_{X^{p,r}} < \infty\}$$

and note that the Schrödinger operator $\Delta + q$ maps $X^{p,r}(\Omega) \rightarrow L^{p'}(\Omega)$ continuously. Throughout this paper we denote by n the unit outer normal to Ω .

DEFINITION 1.1 For a potential $q \in I^\mu(H)$ with $H \cap \text{supp}(q) \subset \text{int}\{\psi = 0\}$, the Cauchy data of the Schrödinger operator $\Delta + q$ relative to $X^{p,r}(\Omega)$ is

$$(1.6) \quad \mathcal{CD}_q = \mathcal{CD}_q^{p,r} = \left\{ \left(u|_{\partial\Omega}, \frac{\partial u}{\partial n} \Big|_{\partial\Omega} \right) : u \in X^{p,r}(\Omega), (\Delta + q)u = 0 \text{ on } \Omega \right\}.$$

By Sobolev embedding, \mathcal{CD}_q is a subspace of $W^{2-\frac{1}{r},r}(\partial\Omega) \times W^{1-\frac{1}{r},r}(\partial\Omega)$. Observe that if the Dirichlet-to-Neumann map Λ_q is defined (on $W^{2-\frac{1}{r},r}(\partial\Omega)$, say), then \mathcal{CD}_q is simply its graph. We will construct certain nontrivial, exponentially growing solutions $u \in X^{p,r}(\Omega)$ so that, for the potentials considered, \mathcal{CD}_q is in fact nontrivial. We can now state our first result.

THEOREM 1.2 Suppose that for $j = 1, 2$, $H_j \Subset \Omega$ are submanifolds of codimension k_j . Suppose further that $q_j \in I^{\mu_j}(H_j)$ are real potentials with $\nu_0(k_j) - k_j < \mu_j < 1 - k_j$ and $\text{supp}(q_j) \Subset \Omega$. Let p and r satisfy

$$2 \leq r < \min(r_0(k_1, \nu_1), r_0(k_2, \nu_2))$$

and

$$\max(p_0(k_1, \nu_1), p_0(k_2, \nu_2)) < p < \infty,$$

and suppose that $\psi \equiv 0$ on a neighborhood of $H_1 \cup H_2$. Then $\mathcal{CD}_{q_1} = \mathcal{CD}_{q_2}$ relative to $X^{p,r}(\Omega)$ implies that $q_1 = q_2$ on Ω .

We also show that under the same assumptions as in Theorem 1.2 for the potential we have a reconstruction procedure; that is, we can reconstruct q from \mathcal{CD}_q (see Theorem 3.1 for more details).

Global uniqueness was established in [28] for $n \geq 3$ (for smooth potentials) and [23] for $n = 2$; for $n \geq 3$ this was extended to $q \in L^\infty$ in [24]. The regularity was further lowered to $q \in L^{n/2}$ in unpublished work of R. Lavine and A. Nachman and to potentials of small norm in the Fefferman-Phong class in [4]. Note that for $-\frac{n-2}{n}k \leq \mu < 0$, $k < \frac{n}{2}$, a general element of $I^\mu(H)$ fails to be in $L^{n/2}(\Omega)$.

The isotropic conductivity problem, where one considers the Dirichlet-to-Neumann map for $L_\gamma = \nabla(\gamma \cdot \nabla)$, can be reduced to the Schrödinger problem via the substitution

$$q = -\frac{\Delta(\gamma^{1/2})}{\gamma^{1/2}},$$

and thus the analogue of the theorem holds for conductivities $\gamma_j \in I^{-k-1-\epsilon}(H_j) \hookrightarrow C^{1+\epsilon}(\overline{\Omega}) \forall 0 < \epsilon < 1$. Currently, the best general global uniqueness result known for $n \geq 3$ is for $\gamma \in C^{3/2}$, proven in [25], building on [2] and using the general argument of [28], while the best result known for $n = 2$ is $\gamma \in W^{1,p}(\Omega)$, $p > 2$, proven in [3] using the $\bar{\partial}$ technique of [1, 22, 23]. Global uniqueness for piecewise-analytic conductivities was proven [18], and special types of jump discontinuities were treated in [15].

Here, we will follow the general argument of [28], although employing a different integral identity so as to avoid difficulties when applying Green's theorem. It is this identity that makes $X^{p,r}$ a convenient space for the problem; indeed, both sides of

$$\int_{\Omega} \Delta u \cdot v - u \cdot \Delta v \, dx = \int_{\partial\Omega} \partial_n u \cdot v - u \cdot \partial_n v \, d\sigma$$

are continuous with respect to $\|\cdot\|_{X^{p,r}}$ and thus Green's theorem holds for $u, v \in X^{p,r}(\Omega)$.

We now start the proof of Theorem 1.2, so as to motivate the technicalities that follow.

PROOF OF THEOREM 1.2: Given a submanifold H of codimension k and a potential $q \in I^\mu(H)$ with $\mu < 1 - k$, we will construct exponentially growing solutions of (1.2) belonging to $X^{p,r}(\Omega)$, of the form $v(x) = e^{\rho \cdot x}(1 + \psi(x, \rho))$, with $\rho \in \mathbb{C}^n$ satisfying $\rho \cdot \rho = 0$.

Let $v_1 \in X^{p,r}(\Omega)$ be a solution of $(\Delta + q_1)v_1 = 0$. By the hypothesis of Theorem 1.2, there is a solution $v_2 \in X^{p,r}(\Omega)$ of $(\Delta + q_2)v_2 = 0$ with

$$(1.7) \quad v_2|_{\partial\Omega} = v_1|_{\partial\Omega} \quad \text{and} \quad \frac{\partial v_2}{\partial n}\Big|_{\partial\Omega} = \frac{\partial v_1}{\partial n}\Big|_{\partial\Omega}.$$

Let $w_2 \in X^{p,r}(\Omega)$ be any other solution to $(\Delta + q_2)w_2 = 0$. Then

$$(\Delta + q_2)(v_1 - w_2) = (\Delta + q_2)v_1 = (\Delta + q_1 + (q_2 - q_1))v_1 = (q_2 - q_1)v_1,$$

so that

$$\begin{aligned}
(1.8) \quad & \int_{\Omega} (q_2 - q_1) v_1 w_2 dx \\
&= \int_{\Omega} (\Delta + q_2)(v_1 - v_2) \cdot w_2 dx \\
&= \int_{\Omega} (\Delta + q_2)(v_1 - v_2) \cdot w_2 - (v_1 - v_2) \cdot (\Delta + q_2) w_2 dx \\
&= \int_{\Omega} \Delta(v_1 - v_2) \cdot w_2 - (v_1 - v_2) \cdot \Delta w_2 dx \\
&= \int_{\partial\Omega} \frac{\partial}{\partial n} (v_1 - v_2) \cdot w_2 - (v_1 - v_2) \cdot \frac{\partial}{\partial n} w_2 d\sigma = 0,
\end{aligned}$$

where the application of Green's theorem is valid since $v_1 - v_2$ and $w_2 \in X^{p,r}$ and the last equality holds by (1.7). If we carry this out for the solutions v_1 and w_2 constructed below for complex frequencies ρ_1 and ρ_2 satisfying $\rho_1 + \rho_2 = -i\xi$, with $\xi \in \mathbb{R}^n \setminus 0$, then we have, as in [28],

$$\begin{aligned}
(1.9) \quad & 0 = \int_{\Omega} (q_1 - q_2) e^{(\rho_1 + \rho_2) \cdot x} (1 + \psi_1(x, \rho_1))(1 + \psi_2(x, \rho_2)) dx \\
&= \widehat{(q_1 - q_2)}(\xi) + \int_{\Omega} e^{-i\xi \cdot x} (q_1 - q_2) (\psi_1 + \psi_2 + \psi_1 \psi_2) dx.
\end{aligned}$$

If one can do this for pairs (ρ_1, ρ_2) with $|\rho_j| \rightarrow \infty$ and show that the last integral $\rightarrow 0$ as $|\rho| \rightarrow \infty$, then $\hat{q}_1(\xi) = \hat{q}_2(\xi)$; doing this for all $\xi \in \mathbb{R}^n$ will finish the proof of Theorem 1.2. \square

As is well-known, $v(x) = e^{\rho \cdot x} (1 + \psi(x))$ is a solution of the Schrödinger equation if and only if ψ is a solution of

$$(1.10) \quad (\Delta_{\rho} + q)\psi = -q(x) \quad \text{where} \quad \Delta_{\rho} = \Delta + 2\rho \cdot \nabla.$$

We will show in Proposition 2.6 that (1.10) is uniquely solvable, with some decay in $|\rho|$, in a Banach space of finite-regularity conormal distributions associated with H , yielding exponentially growing solutions $v_j \in X^{p,r}$ to (1.2) that allow the argument above to be carried out. In Section 3, this result is extended to a hybrid global space, and this is applied to obtain reconstruction of the potential from the Cauchy data, following the general argument of [22]. Finally, in Section 4 we show that uniqueness can fail in a weak formulation of the problem for potentials with very strong singularities on a hypersurface, with blowup rates corresponding to those of distributions conormal of order greater than 1 for H .

2 Uniqueness for Conormal Potentials

As described in the introduction, to prove Theorem 1.2, it suffices to construct exponentially growing solutions to (1.2) of the form $v(x) = e^{\rho \cdot x}(1 + \psi(x, \rho))$ for $\rho \cdot \rho = 0$, $|\rho| \rightarrow \infty$, so that the second integral in (1.9) tends to 0 as $|\rho| \rightarrow \infty$. To do this for potentials $q \in I^\mu(H)$, the standard space of (infinite-regularity) conormal distributions of order μ associated with the codimension k submanifold H , we will also need to formulate Banach spaces of finite-regularity conormal distributions in \mathbb{R}^n associated with H . Rather than working in unnecessary generality, we will restrict ourselves to the spaces needed here; unlike [20, 21], where several other types of finite-regularity conormal spaces are defined using iterated regularity with respect to Lie algebras of tangent vector fields, we impose the finite-regularity assumption directly on the symbols in the oscillatory representations of the distributions, using symbol classes modeled on those of [29].

For $l \in \mathbb{R}$, let C_*^l denote the Zygmund space of order l on \mathbb{R}^n [29]. Thus, if $\psi_0(D) + \sum_{i=1}^{\infty} \psi_i(D) = I$ is a Littlewood-Paley decomposition, with $\psi_i(\xi) = \psi_1(\frac{\xi}{2^{i-1}})$, $i \geq 1$, then

$$\|u\|_{C_*^l} = \sup_i 2^{li} \|\psi_i(D)u\|_{L^\infty(\mathbb{R}^n)}.$$

Recall that if $l \geq 0$, $l \notin \mathbb{Z}$, then $C_*^l = C^{[l], l-[l]}(\mathbb{R}^n)$.

Now fix an order $m \in \mathbb{R}$, an $N \in \mathbb{N}$, and a sequence $\vec{\delta} = (\delta_1, \dots, \delta_N)$ of numbers $0 \leq \delta_j \leq 1$, $1 \leq j \leq N$. For any multi-index $\alpha \in \mathbb{Z}_+^k$, let $\delta(\alpha) = \sum_{j=1}^{|\alpha|} \delta_j$, setting $\delta(0) = 0$ for convenience, and $|\vec{\delta}| = \sum_{j=1}^N \delta_j$.

DEFINITION 2.1

(i) We define

$$C_*^l S^{m, \vec{\delta}}(\mathbb{R}^n \times \mathbb{R}^k) = \{a(x, \theta) : \|\partial_\theta^\alpha a(\cdot, \theta)\|_{C_*^l} \leq C_\alpha (1 + |\theta|)^{m - \delta(\alpha)} \forall |\alpha| \leq N\}$$

and

$$\|a\|_{C_*^l S^{m, \vec{\delta}}} = \max_{0 \leq |\alpha| \leq N} (1 + |\theta|)^{-m + \delta(\alpha)} \|\partial_\theta^\alpha a(\cdot, \theta)\|_{C_*^l}.$$

(ii) If H is a smooth, codimension- k submanifold with compact closure, then $C_*^l I^{m, \vec{\delta}}(H)$ is the space of locally finite sums of distributions of the form $u(x) = \int_{\mathbb{R}^k} e^{iF(x) \cdot \theta} a(x, \theta) d\theta$ with $a \in C_*^l S^{m, \vec{\delta}}$, where $F(x) = (F_1(x), \dots, F_k(x))$ are local defining functions for H .

Remarks. 1. $C_*^l S^{m, \vec{\delta}}$ is a Banach space with respect to the norm defined in (i), and $C_*^l I^{m, \vec{\delta}}(H)$ inherits this structure.

2. $\partial_\theta : C_*^l S^{m, \vec{\delta}} \rightarrow C_*^l S^{m - \delta_1, \vec{\delta}'}$, with $\vec{\delta}' = (\delta_2, \dots, \delta_N)$ and $\partial_x : C_*^l S^{m, \vec{\delta}} \rightarrow C_*^{l-1} S^{m, \vec{\delta}}$ continuously.

3. $C_*^l I^{m, \vec{\delta}}(H)$ is well-defined, since changing the defining functions of H locally corresponds to a change of variable in x , which leaves the symbol class invariant.

4. The usual (infinite-regularity) conormal space $I^m(H)$ has a continuous inclusion with respect to its Fréchet structure: $I^m(H) \hookrightarrow C_*^l I^{m, \vec{\delta}}(H)$ for any l, N , and $\vec{\delta}$.

PROPOSITION 2.2 *Let $H \subset \Omega \Subset \mathbb{R}^n$, with $\text{codim}(H) = k$, $l \notin \mathbb{Z}$ and $\vec{\delta} = (\delta_1, \dots, \delta_N)$.*

(i) *If $l > 1$, $-k \leq m < -\frac{k}{2}$, $N \geq 1$, $\delta_1 > 0$, and $m - |\vec{\delta}| < -k$, then*

$$(2.1) \quad C_*^l I^{m, \vec{\delta}}(H) \hookrightarrow L^p(\Omega) \quad \text{continuously for all } 1 \leq p < \frac{k}{m+k}.$$

(ii) *If $l > 2$, $N \geq 2$, $\delta_1 > 1 - \frac{k}{2}$, and $|\vec{\delta}| > 1$, then for each smooth function \tilde{F} vanishing on H ,*

$$(2.2) \quad v \in C_*^l I^{-k, \vec{\delta}}(H) \implies \tilde{F} \cdot v \in W^{1,p}(\Omega) \quad \text{for } 1 \leq p < \frac{k}{1-\delta_1},$$

and thus, for $G \Subset \overline{\Omega} \setminus H$, $C_^l I^{-k, \vec{\delta}}(H) \hookrightarrow W^{1,p}(G)$ continuously.*

(iii) *Suppose $l > 3$. For $\max(\frac{2}{3}, 1 - \frac{k}{4}) \stackrel{\text{def}}{=} v_0(k) < v < 1$, set*

$$\vec{\delta}_0 = (v, 1-v, 2v-1, 1-v, 1-v, 3v-2).$$

Then, for any \tilde{F} vanishing on H ,

$$(2.3) \quad v \in C_*^l I^{-k, \vec{\delta}_0}(H) \implies \tilde{F}^3 \cdot v \in W^{2,r}(\Omega) \quad \text{for } 1 \leq r < \frac{k}{2(1-v)},$$

and thus, for $G \Subset \overline{\Omega} \setminus H$, $C_^l I^{-k, \vec{\delta}_0}(H) \hookrightarrow W^{2,r}(G)$ continuously.*

PROOF: (i) Since L^p and $C_*^l I^{m, \vec{\delta}}(H)$ are diffeomorphism-invariant, it suffices to assume that, with respect to coordinates $x = (x', x'') \in \mathbb{R}^{n-k} \times \mathbb{R}^k$,

$$H = \{x'' = 0\} \quad \text{and} \quad u(x) = \int_{\mathbb{R}^k} e^{ix'' \cdot \theta} a(x, \theta) d\theta, \quad a \in C_*^l S^{m, \vec{\delta}}.$$

We then have

$$u(x', x'') = \int e^{ix'' \cdot \theta} \left[a(x', 0, \theta) + \sum_{0 < |\alpha''| < [l]} \frac{1}{(\alpha'')!} \partial_{x''}^{\alpha''} a(x', 0, \theta) (x'')^{\alpha''} + R_{[l]}(x', x'', \theta) \right] d\theta,$$

where

$$R_{[l]}(x, \theta) = \sum_{|\alpha''|=[l]} b_{\alpha''}(x, \theta) (x'')^{\alpha''} \quad \text{with } |b_{\alpha''}(x, \theta)| \leq C(1+|\theta|)^m \forall \alpha''.$$

Since $a(x', 0, \cdot) \in L^q(\mathbb{R}_{\theta}^k) \forall q > -\frac{k}{m}$ and $-\frac{k}{m} < 2$, the Hausdorff-Young inequality implies that $\int e^{ix'' \cdot \theta} a(x', 0, \theta) d\theta \in L^{q'}(\mathbb{R}_{x''}^k) \forall 2 \leq q' < \frac{k}{m+k}$ uniformly

in x' , and thus belongs to $L^p(\Omega)$ for all $2 \leq p < \frac{k}{m+k}$; since it is compactly supported, the range is in fact $1 \leq p < \frac{k}{m+k}$. For the second term, note that for each function $a_{\alpha''}(x', \theta) \stackrel{\text{def}}{=} \frac{1}{(\alpha'')!} \partial^{\alpha''} a(x', 0, \theta)$,

$$\begin{aligned} \int e^{ix'' \cdot \theta} a_{\alpha''}(x', \theta) (x'')^{\alpha''} d\theta &= \int \left(\frac{1}{i} \partial_{\theta} \right)^{\alpha''} (e^{ix'' \cdot \theta}) a_{\alpha''}(x, \theta) d\theta \\ &= \int e^{ix'' \cdot \theta} \left(\frac{-1}{i} \partial_{\theta} \right)^{\alpha''} (a_{\alpha''}(x, \theta)) d\theta, \end{aligned}$$

whose amplitude is $\leq C(1 + |\theta|)^{m-\delta(\alpha'')}$ and hence is treated by the same argument as for $a_0 = a(x', 0, \theta)$.

In the final term, we integrate by parts:

$$\int e^{ix'' \cdot \theta} b_{\alpha''}(x, \theta) (x'')^{\alpha''} d\theta = \int e^{ix'' \cdot \theta} (i \partial_{\theta})^{\alpha''} b_{\alpha''}(x, \theta) d\theta,$$

and since $|\partial_{\theta}^{\alpha''} b(x, \theta)| \leq C(1 + |\theta|)^{m-|\bar{\delta}|} \in L^1(\mathbb{R}_{\theta}^k)$ uniformly in x' , this yields a bounded function of $x \in \Omega$.

(ii) If $v \in C_*^l I^{-k, \bar{\delta}}(H)$, then $v \in L^p \forall p < \infty$ by part (i). From

$$v(x) = \int e^{ix'' \cdot \theta} a(x, \theta) d\theta, \quad a \in C_*^l S^{-k, \bar{\delta}},$$

one finds

$$\partial_{x_j} v(x) = \int e^{ix'' \cdot \theta} (i \theta_j a + \partial_{x_j} a) d\theta \in C_*^l I^{1-k, \bar{\delta}}(H) + C_*^{l-1} I^{-k, \bar{\delta}}(H)$$

for $n - k + 1 \leq j \leq n$; if $1 \leq j \leq n - k$, only the second term is present. The second term is covered by part (i) and hence is in $L^p \forall p < \infty$. If the first term is multiplied by some x_{j_0} , $n - k + 1 \leq j_0 \leq n$, and integrated by parts, it becomes an element of $C_*^l I^{1-k-\delta_1, \bar{\delta}'}(H)$, with $\bar{\delta}' = (\delta_2, \dots, \delta_N)$, which by (2.1) is in L^p , $1 \leq p < \frac{k}{1-\delta_1}$. Since any \tilde{F} vanishing on H can be represented as a linear combination of x_{j_0} 's with smooth coefficients, $\tilde{F}(x) \nabla v \in L^p$, and so $v \in W^{1,p}(G)$ for any set G on which $|F|$ is bounded below.

(iii) By (i) and (ii) above, both v and $\tilde{F} \cdot \nabla v \in L^r$. Now, arguing as in (ii), for $n - k + 1 \leq j, j' \leq n$, we have

$$\begin{aligned} \partial_{x_j x_{j'}}^2 v &= \int e^{ix'' \cdot \theta} (-\theta_j \theta_{j'} a + i(\theta_j a_{x_{j'}} + \theta_{j'} a_{x_j}) + a_{\theta_j \theta_{j'}}) d\theta \\ &\in C_*^l I^{2-k, \bar{\delta}_0}(H) + C_*^{l-1} I^{1-k, \bar{\delta}_0}(H) + C_*^{l-2} I^{-k, \bar{\delta}_0}(H), \end{aligned}$$

with simpler expressions if one or both of j or j' is $\leq n - k$. By (i), the last term is in $L^r \forall r < \infty$, if $l > 3$, $N \geq 1$. Integrating by parts and using (ii), x_{j_0} times the second term is in L^r , $1 \leq r < \frac{1}{1-\delta_1} = \frac{1}{1-\nu}$ if $l > 2$, $N \geq 2$. As for the first term, $x_{j_0} x_{j_1} x_{j_2}$ times it is seen, after integrating by parts three times, to be in

$C_*^l I^{2-k-(\delta_1+\delta_2+\delta_3), \vec{\delta}_0''}(H) = C_*^l I^{2-k-2\nu, \vec{\delta}_0''}(H)$ for $\vec{\delta}_0'' = (1-\nu, 1-\nu, 3\nu-2)$, which, by (i), $\hookrightarrow L^r(\Omega) \forall 1 \leq r < \frac{k}{2(1-\nu)}$ if $l > 2$, since $-k \leq 2-k-2\nu < -\frac{k}{2}$ if $1 - \frac{k}{4} < \nu \leq 1$ and $2-k-|\vec{\delta}| = 2-3\nu-k < k$ if $\nu > \frac{2}{3}$. Thus, for any \tilde{F} vanishing on H , $\tilde{F}^3 \cdot C_*^l I^{-k, \vec{\delta}_0}(H) \hookrightarrow W^{2,r}(\Omega)$ if $1 \leq r < \frac{k}{2(1-\nu)}$, $l > 3$. \square

We also have the following:

PROPOSITION 2.3 *If $A \in \Psi^r(\mathbb{R}^n)$ is properly supported with $r < 0$, then for any $l, m \in \mathbb{R}$ and any $\vec{\delta}$, and any $\epsilon > 0$,*

$$A : C_*^l I^{m, \vec{\delta}}(H) \rightarrow C_*^l I^{m+r+\epsilon, \vec{\delta}}(H)$$

and $\|A\|$ is bounded by a finite number of seminorms of $\sigma(A)$ in $S_{1,0}^r(\mathbb{R}^n \times (\mathbb{R}^n \setminus 0))$.

PROOF: Write

$$Af(x) = \int e^{i(x-y)\cdot\xi} a(x, y, \xi) f(y) d\xi dy, \quad a \in S_{1,0}^r,$$

and

$$u(y) = \int e^{iF(y)\cdot\theta} b(y, \theta) d\theta, \quad b \in C_*^l S^{m, \vec{\delta}}.$$

Then

$$\begin{aligned} Au(x) &= \int e^{i[(x-y)\cdot\xi + F(y)\cdot\theta]} a(x, y, \xi) b(y, \theta) d\theta d\xi dy \\ &= \int e^{iF(x)\cdot\theta} c(x, \theta) d\theta, \end{aligned}$$

where

$$(2.4) \quad c(x, \theta) = \int e^{i[(x-y)\cdot\xi + (F(y)-F(x))\cdot\theta]} a(x, y, \xi) b(y, \theta) d\xi dy,$$

and we need to show that $c = c[b] \in C_*^l S^{m+r+\epsilon, \vec{\delta}} \forall \epsilon > 0$. Furthermore, since H is compact, we may assume that the symbol $a(x, y, \xi)$ vanishes for x and y outside some compact set.

Let $\langle \theta \rangle = (1 + |\theta|^2)^{1/2}$. Setting $b_i(y, \theta) = \psi_i(D_y)(b(y, \theta))$ and $c_j(x, \theta) = \psi_j(D_x)(c(x, \theta))$, we have

$$\|\partial_\theta^\alpha b_i(\cdot, \theta)\|_{L^\infty} \leq C 2^{-li} \langle \theta \rangle^{m-\delta(\alpha)} \quad \forall |\alpha| \leq N$$

and want to prove that

$$\|\partial_\theta^\alpha c_j(\cdot, \theta)\|_{L^\infty} \leq C_\epsilon 2^{-lj} \langle \theta \rangle^{m+r+\epsilon-\delta(\alpha)} \quad \forall |\alpha| \leq N.$$

We first consider the case of $\alpha = 0$. Let $\{\tilde{\psi}_i\}_{i=0}^\infty$ and $\{\tilde{\tilde{\psi}}_i\}_{i=0}^\infty$ be bounded families in $S_{1,0}^0$ with $\tilde{\psi}_i = 1$ on $\text{supp}(\psi_i)$ and $\tilde{\tilde{\psi}}_i = 1$ on $\text{supp}(\tilde{\psi}_i)$; then one can write $c_j = T_{ij}(b_i)$, where the operator T_{ij} has the Schwartz kernel

$$K_{ij}(x, y; \theta) = \int e^{i[(x-z) \cdot \zeta + (z-w) \cdot \xi + (F(w)-F(z)) \cdot \theta + (w-y) \cdot \eta]} \psi_j(\zeta) a(z, w, \xi) \tilde{\psi}_i(\eta) d\zeta dz d\xi dw d\eta.$$

It suffices to show that $\sum_{i=0}^\infty \|T_{ij}(b_i)\|_{L^\infty} \leq C_\epsilon 2^{-lj} \langle \theta \rangle^{m+r+\epsilon}$.

As in the proof of Proposition 2.2, we may assume that $F(x) = x''$, where $x = (x', x'') \in \mathbb{R}^{n-k} \times \mathbb{R}^k$; for $\theta \in \mathbb{R}^k$, let $\theta^* = (0, \theta) = DF^*(x)(\theta) \forall x$. Thus, the phase function of $K_{ij}(x, y; \theta)$ is

$$\phi = (x-z) \cdot \zeta + (z-w) \cdot \xi + (w''-z'') \cdot \theta + (w-y) \cdot \eta.$$

Let $\chi \in C_0^\infty(\mathbb{R})$, $\chi(t) = 1$, for $|t| \leq \frac{1}{2}$, $\text{supp}(\chi) \subset \{|t| \leq \frac{3}{4}\}$. Write

$$\begin{aligned} K_{ij}(x, y; \theta) &= \int e^{i\phi} \psi_j(\zeta) \tilde{\psi}_i(\eta) \left[\chi\left(\frac{|\xi - \zeta - \theta^*|}{|\theta|}\right) + (1-\chi)\left(\frac{|\xi - \zeta - \theta^*|}{|\theta|}\right) \right] \\ &\quad \times a(z, w, \xi) d\zeta d\eta d\xi dz dw \\ &= K_{ij}^0(x, y; \theta) + K_{ij}^\infty(x, y; \theta) \end{aligned}$$

with $T_{ij} = T_{ij}^0 + T_{ij}^\infty$ the corresponding operator decomposition.

K_{ij}^∞ may be analyzed by noting that $|\xi - \zeta - \theta^*| \geq c \max(|\xi - \zeta|, |\theta|)$ on $\text{supp}(1-\chi)$, and thus, using

$$\left(\frac{\xi - \zeta - \theta^*}{i|\xi - \zeta - \theta^*|^2} \cdot \nabla_z \right) (e^{i\phi}) = e^{i\phi},$$

we may integrate by parts $n+M$ times in z and then integrate in ξ to obtain

$$K_{ij}^\infty(x, y; \theta) = \int e^{i[(x-z) \cdot \zeta + (w''-z'') \cdot \theta + (w-y) \cdot \eta]} \psi_j(\zeta) \tilde{\psi}_i(\eta) A(z, w; \theta) d\zeta d\eta dz dw$$

with $|\partial_z^\alpha \partial_w^\beta A(z, w; \theta)| \leq C_{\alpha, \beta, M} \langle \theta \rangle^{-M} \forall \alpha, \beta \in \mathbb{Z}_+^n, M \in \mathbb{N}$. If $\max(2^i, 2^j) < \frac{1}{4} \langle \theta \rangle$, we then simply integrate in all variables to obtain $|K_{ij}^\infty| \leq c 2^{n(i+j)} \langle \theta \rangle^{-M}$. If $\langle \theta \rangle < \frac{1}{4} \min(2^i, 2^j)$, then $|\zeta + \theta^*| \geq c|\zeta| \geq c2^j$ and $|\eta + \theta^*| \geq c|\eta| \geq c2^i$, so we may first integrate by parts in z and w to obtain $|K_{ij}^\infty| \leq c 2^{-M'(i+j)} \langle \theta \rangle^{-M}$.

In the following, we denote $a \sim b$ if either $a, b < 4$ or $\frac{1}{4}a \leq b \leq 4a$.

If $\langle \theta \rangle \sim 2^i > 2^{j+1}$ or $\langle \theta \rangle \sim 2^j > 2^{i+1}$, we may integrate by parts in z or w alone to obtain $|K_{ij}^\infty| \leq c 2^{-M'i} \langle \theta \rangle^{-M}$ or $|K_{ij}^\infty| \leq c 2^{-M'j} \langle \theta \rangle^{-M}$, respectively, $\forall M'$. If $2^{i+1} < \langle \theta \rangle < 2^{j-1}$, then $|\zeta + \theta^*| \geq c2^j$ and $|\eta + \theta^*| \geq c|\theta^*|$, so integration by parts in z and w gives $|K_{ij}^\infty| \leq c 2^{-M'j} \langle \theta \rangle^{-M}$; similarly, for $2^{j+1} < \langle \theta \rangle < 2^{i-1}$, we obtain $|K_{ij}^\infty| \leq c 2^{-M'i} \langle \theta \rangle^{-M}$. Finally, if $\langle \theta \rangle \sim 2^i \sim 2^j$, we simply integrate

and get $|K_{ij}^\infty| \leq c2^{n(i+j)}\langle\theta\rangle^{-M} \leq c\langle\theta\rangle^{-(M-2n)}$. Using $\|b_i\|_{L^\infty} \leq c2^{-li}\langle\theta\rangle^m$ and summing in i yields $\sum_{i=0}^\infty \|T_{ij}^\infty(b_i)\|_{L^\infty} \leq c2^{-M'j}\langle\theta\rangle^{-M} \forall M, M'$.

On the other hand, K_{ij}^0 may be analyzed using stationary phase in ξ and w : First rewrite

$$(2.5) \quad K_{ij}^0(x, y; \theta) = \int e^{i[(x-z)\cdot\zeta + (z-y)\cdot\eta]} \psi_j(\zeta) \tilde{\psi}_i(\eta) \\ \times \left[\int e^{i|\theta| [w'' \cdot \frac{\theta}{|\theta|} - w \cdot \xi]} |\theta|^n a\left(z, z + w, |\theta| \left(\xi + \frac{\eta}{|\theta|}\right)\right) \right. \\ \left. \times \chi\left(\left|\xi - \frac{\theta^*}{|\theta|} - \sigma\right|\right) d\xi dw \right]_{\sigma = \frac{\xi - \eta}{|\theta|}} dz d\zeta d\eta.$$

The domain of integration for the inner integral contains a critical point only if $|\sigma| < 2$. In this case the unique nondegenerate critical point is $\xi = \frac{\theta^*}{|\theta|}$, $w = 0$. Applying the method stationary phase and analyzing error terms using [14, theorem 7.7.7] for $|\sigma| < 2$, or just using integration by parts for $|\sigma| \geq 2$, we see that the inner integral is

$$a(z, z, \eta + \theta^*) \chi\left(\frac{|\eta - \xi|}{|\theta|}\right) + e_1\left(z, \frac{\eta}{|\theta|}, \frac{\xi}{|\theta|}, \eta + \theta^*\right)$$

with $e_1 \in S_{1,0}^{r-1}((\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n) \times \mathbb{R}^n \setminus 0)$, and thus

$$(2.6) \quad K_{ij}^0(x, y; \theta) = \int e^{i[(x-z)\cdot\zeta + (z-y)\cdot\eta]} \psi_j(\zeta) \tilde{\psi}_i(\eta) \\ \times \left[a(z, z, \eta + \theta^*) \chi\left(\frac{|\eta - \xi|}{|\theta|}\right) \right. \\ \left. + e_1\left(z, \frac{\eta}{|\theta|}, \frac{\xi}{|\theta|}, \eta + \theta^*\right) \right] d\zeta dz d\eta.$$

Note that $S_{1,0}^{r-1}$ seminorms of e_1 are uniformly bounded over compact subsets of $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n$. We observe for the phase that $d_z((x-z)\cdot\zeta + (z-y)\cdot\eta) = -\zeta + \eta$ and $(d_\zeta + d_\eta)((x-z)\cdot\zeta + (z-y)\cdot\eta) = x - y$, and we may use these to integrate by parts, M and M' times, respectively, to obtain, for $i \leq j - 2$,

$$|K_{ij}^0(x, y; \theta)| \leq C \int 2^{-Mj} (1 + 2^i |x - y|)^{-M'} \tilde{\psi}_j(\zeta) \langle\theta\rangle^r |\eta|^{-M'} \tilde{\psi}_i(\eta) d\zeta dz d\eta \\ \leq C 2^{-(M-n)j} (1 + 2^i |x - y|)^{-M'} 2^{-(M'-n)i} \langle\theta\rangle^r.$$

Here, we have used $|\eta - \zeta| \geq \frac{1}{2}|\zeta| \geq 2^{j-1}$ and

$$|(d_\zeta + d_\eta)^{M'}(\psi_j(\zeta) a(z, z, \eta + \theta^*) \tilde{\psi}_i(\eta))| \leq \\ \begin{cases} C |\eta|^{r-M'} \tilde{\psi}_j(\zeta) \tilde{\psi}_i(\eta), & |\theta| \leq c2^i, \\ C \langle\theta\rangle^r |\eta|^{-M'} \tilde{\psi}_j(\zeta) \tilde{\psi}_i(\eta), & |\theta| \geq c2^i. \end{cases}$$

Thus, $\int |K_{ij}^0(x, y)| dy \leq C 2^{-(M-n)j} 2^{-M'i} \langle \theta \rangle^r$, so that

$$\sum_{i=0}^{j-2} \|T_{ij}^0(b_i)\|_{L^\infty} \leq C \sum_{i=0}^{j-2} 2^{-(M-n)j} 2^{-M'i} \langle \theta \rangle^r 2^{-li} \langle \theta \rangle^m \leq C c^{-lj} \langle \theta \rangle^{m+r}$$

for $M > n + l$. Similarly, for $i \geq j + 2$, we obtain the estimate

$$|K_{ij}^0(x, y)| \leq C 2^{-(M-n)i} (1 + 2^j |x - y|)^{-M'} 2^{-(M'-n)j} \langle \theta \rangle^r,$$

leading to

$$\sum_{i=j+2}^{\infty} \|T_{ij}^0(b_i)\|_{L^\infty} \leq C \sum_{i=j+2}^{\infty} 2^{-M'j} 2^{-(M-n)i} \langle \theta \rangle^r 2^{-li} \langle \theta \rangle^m \leq C 2^{-lj} \langle \theta \rangle^{m+r}$$

for $M' > l$, $M > n - l$.

To analyze the contributions from $|i - j| \leq 1$, we perform in (2.6) an additional stationary phase in z and ζ , with critical point $z = x$ and $\zeta = \eta$. Rewriting the integral as in formula (2.5), we see that

$$K_{ij}^0(x, y) = \int e^{i(x-y)\cdot\eta} \psi_j(\eta) \left[a(x, x, \eta + \theta^*) + e_2\left(x, \frac{\eta}{|\theta|}, \frac{\eta}{|\theta|}, \eta + \theta^*\right) \right] \tilde{\psi}_i(\eta) d\eta$$

with $e_2 \in S_{1,0}^{r-1}((\mathbb{R}^n \times \mathbb{R}^n) \times \mathbb{R}^n \setminus 0)$. Since ψ_j are uniformly bounded in $S_{1,0}^0$, the $S_{1,0}^{r-1}$ seminorms of e_2 are uniformly bounded over compact subsets of $\mathbb{R}^n \times \mathbb{R}^n$. Then

$$K_{ij}^0(x, y) = e^{-i\theta^*\cdot x} \left[\int e^{i(x-y)\cdot\eta} [\psi_j(\eta - \theta^*) h(x, \eta; \theta) \tilde{\psi}_i(\eta - \theta^*)] \tilde{\psi}_i(\eta - \theta^*) d\eta \right] e^{i\theta^*\cdot y}$$

where $h(\cdot, \cdot; \theta) \in S_{1,0}^r(\mathbb{R}^n \times \mathbb{R}^n \setminus 0)$,

$$h(x, \eta; \theta) = a(x, x, \eta) + e_2\left(x, \frac{\eta - \theta^*}{|\theta|}, \frac{\eta - \theta^*}{|\theta|}, \eta\right),$$

is a symbol depending on the parameter θ , $|\theta| > 1$.

Note that multiplication by the exponentials does not affect the $L^\infty \rightarrow L^\infty$ operator norm, while

$$\psi_j(\eta - \theta^*) h(x, \eta; \theta) \tilde{\psi}_i(\eta - \theta^*) \in \begin{cases} S_{1,0}^r(\mathbb{R}^n \times \mathbb{R}^n \setminus 0), & |\theta| \leq c 2^j, \\ \langle \theta \rangle^r \cdot S_{1,0}^0(\mathbb{R}^n \times \mathbb{R}^n \setminus 0), & |\theta| \geq c 2^j, \end{cases}$$

uniformly in θ . Acting on functions with Fourier transforms supported in $B(0, R)$, pseudodifferential operators of order 0 are bounded on L^∞ with norm $\leq c \log R$ [29]. Hence, for $|i - j| \leq 1$,

$$\|T_{ij}(b_i)\|_{L^\infty} \leq \begin{cases} C j 2^{rj} 2^{-lj} \langle \theta \rangle^m \leq C_\epsilon 2^{-lj} \langle \theta \rangle^{m+r+\epsilon}, & |\theta| \leq c 2^j, \\ C j 2^{-lj} \langle \theta \rangle^{m+r} \leq C_\epsilon 2^{-lj} \langle \theta \rangle^{m+r+\epsilon}, & |\theta| \geq c 2^j, \end{cases}$$

as desired.

To handle the derivatives $\partial_\theta^\alpha c$, note first that the application of ∂_θ to the right-hand side of (2.4) yields two terms. Since $\partial_\theta b(x, \theta) \in C_*^l S^{m-\delta_1, (\delta_2, \dots)}$, the oscillatory integral with amplitude $a(x, y, \xi) \partial_\theta b(x, \theta)$ has C_*^l norm $\leq C \langle \theta \rangle^{m+r-\delta_1}$, as desired. On the other hand, because $F(x) = x''$ here, we see that if the derivative hits the phase, then the amplitude becomes $i(F(y) - F(x)) \cdot a \cdot b = i(y'' - x'') \cdot a \cdot b$. Writing $y'' - x'' = -\partial_{\xi''}[(x - y) \cdot \xi + (F(y) - F(x)) \cdot \theta]$, we may then integrate by parts in ξ . The resulting amplitude, $i \partial_\xi a \cdot b$, gives a term whose C_*^l norm is $\leq C \langle \theta \rangle^{m+r-1}$. Higher derivatives $\partial_\theta^\alpha c$, $|\alpha| \leq N$, are handled similarly. \square

To obtain the boundedness of M_q on some of the finite-regularity conormal spaces, we will need the following (similar results are in [13, 26]):

LEMMA 2.4 *Let $a, b \in C(\mathbb{R}^n \times \mathbb{R}^k)$ satisfy*

$$\|a(\cdot, \theta)\|_{C_*^l} \leq C \langle \theta \rangle^m \quad \text{and} \quad \|b(\cdot, \theta)\|_{C_*^{l'}} \leq C \langle \theta \rangle^{m'}$$

*with $l, l' > 0$, $l, l' \notin \mathbb{N}$, and $m + m' < -k$. Then the partial convolution $a *_{\mathbb{R}^k} b$ satisfies*

$$\|a *_{\mathbb{R}^k} b(\cdot, \theta)\|_{C_*^{l''}} \leq C \langle \theta \rangle^{m''},$$

with $l'' = \min(l, l')$ and $m'' = \max((m + k)_+ + m', m + (m' + k)_+, m + m' + k)$. Here, $t_+ = t_+ = \max(t, 0)$ if $t \neq 0$ and $t_+ = \epsilon$ for $t = 0$, with $\epsilon > 0$ arbitrarily small.

SKETCH OF PROOF: This follows easily by decomposing

$$\begin{aligned} (a *_{\mathbb{R}^k} b)(x, \theta) &= \int_{\mathbb{R}^k} a(x, \sigma) b(x, \theta - \sigma) d\sigma \\ &= \left(\int_{|\sigma| \leq \frac{|\theta|}{2}} + \int_{|\theta - \sigma| \leq \frac{|\theta|}{2}} + \int_{\substack{|\sigma| \geq \frac{|\theta|}{2} \\ |\theta - \sigma| \geq \frac{|\theta|}{2}}} \right) a(x, \sigma) b(x, \theta - \sigma) d\sigma, \end{aligned}$$

and estimating each of the three terms using the fact that $C_*^l \times C_*^{l'} \hookrightarrow C_*^{l''}$ continuously. \square

This is then used in the proof of the following:

PROPOSITION 2.5 *Let $H \subset \mathbb{R}^N$ be codimension k . For $q \in I^\mu(H)$, let M_q denote multiplication by $q(x)$. Suppose $\frac{2}{3} - k \leq \mu < 1 - k$. Then, for any $l > 0$, $l \notin \mathbb{N}$,*

$$(2.7) \quad M_q : C_*^l I^{-k, \vec{\delta}_0}(H) \rightarrow C_*^l I^{\tilde{\mu}, \vec{\delta}_0}(H) \quad \text{continuously}$$

where, if we write $\mu = \nu - k$ with $\frac{2}{3} \leq \nu < 1$, $\vec{\delta}_0$ is as in Proposition 2.2(iii):

$$(2.8) \quad \vec{\delta}_0 = (\nu, 1 - \nu, 2\nu - 1, 1 - \nu, 1 - \nu, 3\nu - 2)$$

and $\tilde{\mu} = \mu + \epsilon$ for any $0 < \epsilon < -\mu$.

PROOF: As before, we can assume that for any $l' > 0$, $H = \{x'' = 0\}$,

$$u(x) = \int_{\mathbb{R}^k} e^{ix'' \cdot \theta} b(x, \theta) d\theta, \quad b \in C_*^l I^{-k, \bar{\delta}_0}(H),$$

$$q(x) = \int_{\mathbb{R}^k} e^{ix'' \cdot \theta} a(x, \theta) d\theta, \quad a \in S_{1,0}^\mu \hookrightarrow C^{l'} S^{\mu, (1,1,\dots)},$$

Hence, $M_q u(x) = \int e^{ix'' \cdot \theta} (a * b)(x, \theta) d\theta$, where $*$ denotes the k -dimensional convolution in the θ -variable. By Lemma 2.4,

$$\|a * b(x, \theta)\|_{C_*^l} \leq C \langle \theta \rangle^{\max(\mu, \mu+\epsilon, \mu)} = C \langle \theta \rangle^{\mu+\epsilon} \quad \forall \epsilon > 0.$$

Since $\partial_\theta a \in C_*^{l'} S^{\mu-1, (1,1,\dots)}$ and $\mu - 1 < -k$,

$$\|\partial_\theta(a * b)(x, \theta)\|_{C_*^l} = \|(\partial_\theta a) * b(x, \theta)\|_{C_*^l} \leq C \langle \theta \rangle^{\max(-k, \mu-1+\epsilon, \mu-1)} = C \langle \theta \rangle^{-k}$$

(for $\epsilon < -\mu$), which gives a gain of $\geq \mu - (-k) = \nu$, consistent with $\delta_1 = \nu$. (The additional gain of ϵ we choose to ignore.) By Remark 2 above, noting that $\partial_\theta b \in C_*^{l'} S^{-k-\nu, (1-\nu, 2\nu-1, \dots, 3\nu-2)}$, we have

$$\|\partial_\theta^2(a * b)\|_{C_*^l} = \|(\partial_\theta^2 a) * (\partial_\theta b)\|_{C_*^l} \leq C \langle \theta \rangle^{\max(-k-\nu, \nu-k-1, -k-1)} = C \langle \theta \rangle^{\nu-k-1}$$

since $\nu \geq \frac{1}{2}$, which is a gain of $\delta_2 = -k - (\nu - k - 1) = 1 - \nu$. Since $\partial_\theta^2 a \in C_*^{l'} S^{\mu-2, (1,1,\dots)}$ and $\partial_\theta b$ is as noted above,

$$\|\partial_\theta^3(a * b)\|_{C_*^l} = \|(\partial_\theta^3 a) * (\partial_\theta b)\|_{C_*^l} \leq C \langle \theta \rangle^{\max(-k-\nu, \nu-k-2)} = C \langle \theta \rangle^{-k-\nu},$$

which gives a gain of $\delta_3 = \nu - k - 1 - (-k - \nu) = 2\nu - 1$. Since $\partial_\theta^2 a \in C_*^{l'} S^{\mu-2, (1,1,\dots)}$ and $\partial_\theta^2 b \in C_*^{l'} S^{-k-1, (2\nu-1, 1-\nu, \dots)}$,

$$\|\partial_\theta^4(a * b)\|_{C_*^l} = \|(\partial_\theta^4 a) * (\partial_\theta^2 b)\|_{C_*^l} \leq C \langle \theta \rangle^{\max(-k-1, \nu-k-2, 2\nu-k-3)} = C \langle \theta \rangle^{-k-1},$$

which gives a gain of $\delta_4 = -k - \nu - (-k - 1) = 1 - \nu$. Continuing in this fashion, we may estimate

$$\|\partial_\theta^5(a * b)\|_{C_*^l} = \|(\partial_\theta^5 a) * (\partial_\theta^3 b)\|_{C_*^l} \leq C \langle \theta \rangle^{\nu-k-2},$$

which is consistent with $\delta_5 = 1 - \nu$ if $\nu \geq \frac{2}{3}$, and

$$\|\partial_\theta^6(a * b)\|_{C_*^l} = \|(\partial_\theta^6 a) * (\partial_\theta^3 b)\|_{C_*^l} \leq C \langle \theta \rangle^{-2\nu-k},$$

which is consistent with $\delta_6 = 3\nu - 2$. The x -derivatives, lowering the Zygmund space index and not involving any gain in $\langle \theta \rangle$, are handled in the obvious way. Hence, $a * b \in C_*^{l'} S^{\mu+\epsilon, \bar{\delta}_0}$. \square

Now recall some facts concerning the Faddeev Green's function G_ρ [10]. As is well-known (see, e.g., [28], where this is used implicitly), the families $\{|\rho|G_\rho : \rho \cdot \rho = 0\}$ and $\{G_\rho : \rho \cdot \rho = 0\}$ are uniformly bounded in $\Psi^0(K)$ and $\Psi^{-1}(K)$, respectively, for $K \Subset \mathbb{R}^n$ and hence interpolation implies

$$(2.9) \quad \{|\rho|^{1-t} G_\rho : \rho \cdot \rho = 0\} \subset \Psi^{-t}(K) \quad \text{is bounded} \quad \forall t \in [0, 1].$$

We can now state a local analogue for the finite-regularity conormal spaces of the result of [28] concerning solvability of inhomogeneous equations involving $\Delta_\rho + q(x)$ in weighted L^2 spaces.

PROPOSITION 2.6 *If $q \in I^\mu(H)$ with $\frac{2}{3} - k \leq \mu < 1 - k$, $l > 3$, $l \notin \mathbb{N}$, and $0 \leq \sigma \leq 1$, then for $\vec{\delta}_0$ as in (2.8), the inhomogeneous equation*

$$(2.10) \quad (\Delta_\rho + q)w = g \in C_*^l I^{1-k-\sigma, \vec{\delta}_0}(H)$$

has, for $|\rho|$ large, a unique solution $w \in C_^l I^{-k, \vec{\delta}_0}(H)$, with $\|w\| \leq \frac{C}{|\rho|^\sigma} \|g\|$.*

PROOF: Applying G_ρ to both sides of (2.10), and using (2.9) for $t = 1 - \sigma$ and Proposition 2.3, we are reduced to showing that

$$(I + G_\rho M_q)w = G_\rho g \in C_*^l I^{-k, \vec{\delta}_0}(H)$$

has a unique solution for $|\rho|$ sufficiently large, with $\|w\| \leq C \|G_\rho g\|$. By Proposition 2.5, $M_q : C_*^l I^{-k, \vec{\delta}_0}(H) \rightarrow C_*^l I^{\mu+\epsilon, \vec{\delta}_0}(H)$ for any $0 < \epsilon < -\mu$. Note that $t = \mu + \epsilon - (-k) < 1$, so we can use Proposition 2.5 and (2.9) with this value of t to obtain

$$C_*^l I^{-k, \vec{\delta}_0}(H) \xrightarrow{M_q} C_*^l I^{\mu+\epsilon, \vec{\delta}_0}(H) \xrightarrow{G_\rho} C_*^l I^{-k, \vec{\delta}_0}(H)$$

with norm $\leq C(q)|\rho|^{t-1+\epsilon'} \rightarrow 0$ as $|\rho| \rightarrow \infty$. Hence, for $|\rho|$ sufficiently large, $\|G_\rho M_q\| < \frac{1}{2}$ and $I + G_\rho M_q$ is invertible on $C_*^l I^{-k, \vec{\delta}_0}(H)$. \square

COMPLETION OF THE PROOF OF THEOREM 1.2: Construct two solutions ψ_j , $j = 1, 2$, to (1.10) for potentials $q_j \in I^{\mu_j}(H_j)$. Note that $-q_j$, the right-hand side of (1.10), belongs to $I^{\mu_j}(H) \hookrightarrow C_*^l I^{1-k_j-\sigma_j, \vec{\delta}}(H)$, with $\sigma_j > 0$ since $\mu_j < 1 - k_j$. Thus, we may apply Proposition 2.6 with $g = -q_j$, $j = 1, 2$, and then form as above the corresponding solutions $v_1(x, \rho_1) = e^{\rho_1 \cdot x}(1 + \psi_1(x, \rho_1))$ of $(\Delta + q_1)v_1 = 0$ and $w_2(x, \rho_2) = e^{\rho_2 \cdot x}(1 + \psi_2(x, \rho_2))$ of $(\Delta + q_2)w_2 = 0$, with $\|\psi_j\|_{C_*^l I^{-k_j, \vec{\delta}}} \leq C|\rho|^{-\sigma_j}$. The solutions v_1 and w_2 belong to $X^{p,r}(\Omega)$, with p and r as in (1.3). In fact, each is in $L^p(\Omega)$ and in $W^{2,r}(\Omega)$ away from H by Proposition 2.2(i) and (iii), respectively, since $r < \frac{1}{2(1-\nu)}$. Furthermore, since $q_1 \in L^{k_1/\nu_1-\epsilon} \forall \epsilon > 0$, we have $\Delta v_1 = -q_1 v_1 \in (L^{k_1/\nu_1-\epsilon}) \times (L^s) \forall \epsilon > 0, \forall s < \infty$. Since $p > \frac{k_1}{k_1-\nu_1} \implies p' < \frac{k_1}{\nu_1}$, we thus have $\Delta v_1 \in L^{p'}(\Omega)$, and similarly for w_2 . These solutions are constructed for all large $|\rho|$. Since $n \geq 3$, for any $\xi \in \mathbb{R}^n \setminus 0$ and $\lambda \geq c|\xi|$, one can find $\rho_1, \rho_2 \in \{\rho \cdot \rho = 0\}$ with $|\rho_1| \simeq |\rho_2| \geq \lambda$ and $\rho_1 + \rho_2 = -i\xi$.

By the assumption that $\mathcal{CD}_{q_2} = \mathcal{CD}_{q_1}$, there exists a $v_2 \in X^{p,r}(\Omega)$ such that (1.7) holds. Applying (1.8) and (1.9), it suffices to show that

$$(2.11) \quad \int_{\Omega} e^{-i\xi \cdot x} (q_1 - q_2)(\psi_1 + \psi_2 + \psi_1 \psi_2) dx \rightarrow 0 \quad \text{as } \lambda \rightarrow \infty.$$

Since each $q_j \in L^{k_j/v_j-\epsilon}(\Omega) \forall \epsilon > 0$, and, by (1.3), $p > (\frac{k_j}{v_j})'$, $j = 1, 2$, we have

$$\int_{\Omega} |q_1 - q_2| \cdot |\psi_j| dx \leq \frac{c}{\lambda^{\epsilon_j}}, \quad \epsilon_j > 0,$$

as $\lambda \rightarrow \infty$ by Hölder's inequality. In fact, $p > 2(\frac{k_j}{v_j})'$ by (1.3), so we have

$$\int_{\Omega} |q_1 - q_2| \cdot |\psi_1| \cdot |\psi_2| dx \leq \frac{c}{\lambda^{\epsilon_{12}}}, \quad \epsilon_{12} > 0,$$

as well. Thus, we have shown that $\mathcal{CD}_{q_1} = \mathcal{CD}_{q_2}$ implies that $\widehat{q_1 - q_2}(\xi) = 0 \forall \xi$, and hence $q_1 = q_2$, concluding the proof of Theorem 1.2. \square

3 Reconstruction of the Potential

In this section we prove that the potential q can be obtained constructively from the Cauchy data of $\Delta + q$. We follow here the general method of [22]; see also [17, 23]. However, there are some additional difficulties in our case because we deal with the set of Cauchy data instead of the Dirichlet-to-Neumann map. Moreover, we work with more complicated function spaces due to the singularities of the potential. We will show the following:

THEOREM 3.1 *Suppose that Ω is Lipschitz and $H \Subset \Omega$ is a submanifold of codimension k . Suppose further that $q \in I^{\mu}(H)$ is a real potential with $v_0(k) - k < \mu < 1 - k$ and $\text{supp}(q) \Subset \Omega$. Let p and r satisfy $2 \leq r < r_0(k, v)$ and $\max(p_0(k, v), -\frac{k}{\mu}) < p < \infty$. Then q can be reconstructed on Ω from the Cauchy data \mathcal{CD}_q of $\Delta + q$ on $X^{p,r}(\Omega)$.*

PROOF: To start the discussion of reconstruction, we first show how to obtain a global analogue of Proposition 2.6. For $s \in \mathbb{R}$ and $-1 < \delta < 0$, let $W_{\delta}^{s,2}(\mathbb{R}^n)$ be the weighted Sobolev space denoted by H_{δ}^s in [28], where it is shown that

$$(3.1) \quad \|G_{\rho} f\|_{W_{\delta}^{s+t,2}} \leq \frac{C}{|\rho|^{1-t}} \|f\|_{W_{\delta+1}^{s,2}} \quad \text{for } 0 \leq t \leq 1.$$

Now let $\text{supp}(q) \subset \Omega' \Subset \Omega$ and let $\chi_0 + \chi_{\infty} \equiv 1$ be a partition of unity subordinate to the open cover $\Omega \cup (\Omega')^c = \mathbb{R}^n$. Fix $\frac{2}{3} - k < \mu < 1 - k$, $l > 3$, and let $\tilde{\delta}_0$ be as in (2.8). For $m \leq 1 - k$ and $s \leq -2$, define

$$(3.2) \quad \|f\|_{Y_{\delta}^{m,s}(\mathbb{R}^n)} = \|\chi_0 \cdot f\|_{C_*^l I^{m,\tilde{\delta}_0}(H)} + \|\chi_{\infty} \cdot f\|_{W_{\delta}^{s,2}(\mathbb{R}^n)}.$$

For $m \leq -k$, elements of $I^m(H)$ are in $L_{\text{loc}}^p(\mathbb{R}^n) \forall p < \infty$, and hence $I_{\text{comp}}^m(H) \hookrightarrow W_{\delta}^{s,2}(\mathbb{R}^n) \forall m \leq 1 - k, s \leq -1$. Combining (3.1) with (2.9) and Proposition 2.3, we have, for $0 \leq t \leq 1$,

$$(3.3) \quad \|G_{\rho} f\|_{Y_{\delta}^{m-t,s+t}(\mathbb{R}^n)} \leq \frac{C}{|\rho|^{1-t}} \|f\|_{Y_{\delta+1}^{m,s}(\mathbb{R}^n)},$$

$$m \leq 1 - k - t, \quad s \leq -1 - t, \quad -1 < \delta < 0.$$

Finally, since $\text{supp}(q) \Subset \Omega$, it follows from (2.7) that

$$(3.4) \quad M_q : Y_\delta^{-k,s} \rightarrow Y_\delta^{\mu+\epsilon,s} \quad \forall \epsilon > 0.$$

Arguing as in the proof of Proposition 2.6, but replacing the local finite-regularity spaces with the global spaces $Y_\delta^{m,s}(\mathbb{R}^n)$, we obtain the following result:

PROPOSITION 3.2 *If $q \in I^\mu(H)$ with $\frac{2}{3} - k < \mu < 1 - k$, $s \leq -2$, $-1 < \delta < 0$, and $0 \leq \sigma \leq 1$, then the inhomogeneous equation*

$$(3.5) \quad (\Delta_\rho + q)w = g \in Y_{\delta+1}^{1-k-\sigma,s}(\mathbb{R}^n)$$

has, for $|\rho|$ large, a unique solution $w \in Y_\delta^{-k,s+1-\sigma}(\mathbb{R}^n)$ with $\|w\| \leq \frac{C}{|\rho|^\sigma} \|g\|$.

Next we will construct the boundary values of the exponentially growing solutions on $\partial\Omega$. For this purpose we use the Green's function $G_\rho^q(x, y)$ defined by

$$(3.6) \quad (\Delta + q)G_\rho^q(\cdot, y) = \delta_y \text{ in } \mathbb{R}^n, \quad e_\rho(\cdot)G_\rho^q(\cdot, y) \in Y_\delta^{-k,s}(\mathbb{R}^n),$$

where $y \in \mathbb{R}^n \setminus \overline{\Omega}$, $e_\rho(x) = \exp(-\rho \cdot x)$, and $s < -n/2$. When $|\rho|$ is large enough, equation (3.6) has a unique solution by Proposition 3.2. Next we consider the case when ρ is fixed and sufficiently large.

As $\text{supp}(q) \Subset \Omega$, we see that $\partial\Omega$ has a neighborhood V such that in $V \times V$ Green's function $G_\rho^q(x, y)$ has the same singularities as the Green's function G_0^0 (for the zero potential and $\rho = 0$), that is, $G_\rho^q(x, y) - G_0^0(x, y) \in C^\infty(V \times V)$.

Using the Green's function (3.6), we define the corresponding single- and double-layer potentials

$$S_q\phi(y) = \int_{\partial\Omega} G_\rho^q(x, y)\phi(x)dS_x,$$

$$K_q\phi(y) = \int_{\partial\Omega} \left(\frac{\partial}{\partial n(x)} G_\rho^q(x, y) \right) \phi(x)dS_x, \quad y \notin \partial\Omega,$$

which define continuous operators $S_q : W^{1-\frac{1}{r},r}(\partial\Omega) \rightarrow X|_\Omega \oplus W_{\text{loc}}^{2,r}(\mathbb{R}^n \setminus \Omega)$ and $K_q : W^{2-\frac{1}{r},r}(\partial\Omega) \rightarrow X|_\Omega \oplus W_{\text{loc}}^{2,r}(\mathbb{R}^n \setminus \Omega)$. Here $X \subset \mathcal{D}'(\mathbb{R}^n)$ is the space with the norm $\|\chi_0 \cdot f\|_{C_*^{l,m,\tilde{\delta}_0}(H)} + \|\chi_\infty \cdot f\|_{W_\delta^{2,r}(\mathbb{R}^n)}$. These layer potentials can be considered as operators on the boundary $\partial\Omega$, defined in the principal value sense. Since $\partial\Omega$ is Lipschitz, it follows from the results of [7] that these operators are continuous, $S_q^{\partial\Omega} : W^{1-\frac{1}{r},r}(\partial\Omega) \rightarrow W^{2-\frac{1}{r},r}(\partial\Omega)$ and $K_q^{\partial\Omega} : W^{2-\frac{1}{r},r}(\partial\Omega) \rightarrow W^{2-\frac{1}{r},r}(\partial\Omega)$. Similarly, on $\partial\Omega$ we define normal derivatives of the layer potentials,

$$\partial_n S_q^{\partial\Omega}\phi(y) = \text{P.V.} \int_{\partial\Omega} \left(\frac{\partial}{\partial n(y)} G_\rho^q(x, y) \right) \phi(x)dS_x,$$

$$\partial_n K_q^{\partial\Omega}\phi(y) = \text{P.V.} \int_{\partial\Omega} \left(\frac{\partial}{\partial n(y)} \frac{\partial}{\partial n(x)} G_\rho^q(x, y) \right) \phi(x)dS_x,$$

which are continuous operators $\partial_n S_q^{\partial\Omega} : W^{1-\frac{1}{r},r}(\partial\Omega) \rightarrow W^{1-\frac{1}{r},r}(\partial\Omega)$ and $\partial_n K_q^{\partial\Omega} : W^{2-\frac{1}{r},r}(\partial\Omega) \rightarrow W^{1-\frac{1}{r},r}(\partial\Omega)$.

Next we consider the Calderón projector [5]. We start with the operator

$$A_q(\phi, \psi) = \left(-S_q^{\partial\Omega}\phi + \left(-\frac{1}{2} + K_q^{\partial\Omega} \right) \psi, -\left(\frac{1}{2} + \partial_n S_q^{\partial\Omega} \right) \phi + \partial_n K_q^{\partial\Omega} \psi \right).$$

PROPOSITION 3.3 *Let $Z = W^{2-\frac{1}{r},r}(\partial\Omega) \times W^{1-\frac{1}{r},r}(\partial\Omega)$. Then the operator*

$$A_q : Z / \ker(A_q) \rightarrow Z$$

is semi-Fredholm. Moreover, $-A_q : Z \rightarrow Z$ is a projection operator with range \mathcal{CD}_q and kernel independent of q . In particular, \mathcal{CD}_q is a closed subspace of Z .

PROOF: First we show that the kernel of A_q does not depend on q . Assume that $(\phi, \psi) \in Z$. We consider the function

$$u_{\phi,\psi}(y) = -S_q(\phi) + K_q(\psi), \quad y \in \mathbb{R}^n \setminus \partial\Omega,$$

and the trace operators

$$T_+ : W^{2,r}(\mathbb{R}^n \setminus \Omega) \rightarrow W^{2-\frac{1}{r},r}(\partial\Omega) \times W^{1-\frac{1}{r},r}(\partial\Omega),$$

$$T_- : W^{2,r}(\Omega) \rightarrow W^{2-\frac{1}{r},r}(\partial\Omega) \times W^{1-\frac{1}{r},r}(\partial\Omega),$$

defined by $T_{\pm}u = (u|_{\partial\Omega}, \partial_n u|_{\partial\Omega})$. As the Green's functions $G_{\rho}^q(x, y)$ have the same singularities near $\partial\Omega \times \partial\Omega$ as the standard Green's function of \mathbb{R}^n , we can use the standard jump relations for layer potentials (see, e.g., [8]). We conclude that

$$T_- u_{\phi,\psi} = A_q(\phi, \psi), \quad T_+ u_{\phi,\psi} = (\phi, \psi) + A_q(\phi, \psi).$$

Thus we get that $u = u_{\phi,\psi} \in (e_{\rho})^{-1} Y_{\delta}^{-k,s}$, and it is the unique solution of

$$(3.7) \quad (\Delta + q)u = g_{\phi,\psi} = \psi \delta_{\partial\Omega} + \nabla \cdot (n \phi \delta_{\partial\Omega}) \quad \text{in } \mathbb{R}^n$$

satisfying $e_{\rho}(\cdot)u \in Y_{\delta}^{-k,s}(\mathbb{R}^n)$.

Now, if $(\phi, \psi) \in \ker(A_q)$, we have that

$$(\phi, \psi) = (\phi, \psi) + T_- u_{\phi,\psi} = T_+ u_{\phi,\psi}.$$

Thus, $v = u_{\phi,\psi}$ is the solution of the scattering problem

$$(3.8) \quad \Delta v = 0 \text{ in } \mathbb{R}^n \setminus \Omega, \quad T_+ v = (\phi, \psi), \quad e_{\rho}(\cdot)v \in W_{\delta}^{s,2}(\mathbb{R}^n \setminus \Omega).$$

On other hand, assume that (3.8) has a solution, and let v_0 be the zero continuation of v , that is, $v_0|_{\mathbb{R}^n \setminus \Omega} = v$, $v|_{\Omega} = 0$. Then we conclude that v_0 is a solution of the problem (3.7), and because this solution is unique, we see that $v_0 = u_{\phi,\psi}$. This shows that $(\phi, \psi) \in \ker(A_q)$ if and only if problem (3.8) has a solution. This is obviously independent of q , and thus we see that

$$(3.9) \quad T_- + (e_{\rho} + u_{\phi,\psi}) = (\phi_{\rho} + \phi, \psi_{\rho} + \psi) + A_q(\phi, \psi) \in \text{ran}(A_q).$$

Applying the projection $I + A_q$ to both sides of (3.9) and using $A_q(\phi, \psi) = 0$, we see that

$$0 = (I + A_q)(\phi_\rho + \phi, \psi_\rho + \psi) = (I + A_q)(\phi_\rho, \psi_\rho) + (\phi, \psi).$$

Because A_q and (ϕ_ρ, ψ_ρ) are known, we can thus determine (ϕ, ψ) and the Cauchy data of $v(x)$ on $\partial\Omega$.

Next we consider the range of A_q . A standard application of Green's formula (see, e.g., [8, theorem 3.1]) shows that if $v \in X^{p,r}$ satisfies

$$(\Delta + q)v = 0 \quad \text{in } \Omega$$

and $(\phi, \psi) = T_-v$, then $v = -u_{\phi,\psi}$. (Observe the negative sign, which is due to the fact that we use the exterior normal vector n .) Also, for $(\phi, \psi) \in Z$, we have $\chi_0 u_{\phi,\psi} \in C_*^l I^{-k, \vec{\delta}_0}(H) \subset L^1(\Omega)$ for any $t_1 < \infty$ by Proposition 2.2. As $q \in I^\mu(H) \subset L^2(\Omega)$ for $1 < t_2 < \frac{k}{k+\mu}$, we have $\chi_0 \Delta u_{\phi,\psi} = -\chi_0 q u_{\phi,\psi} \in L^{p'}(\Omega)$ for $\frac{1}{p} < 1 - \frac{1}{t_2}$, i.e., $p > -\frac{k}{\mu}$. Hence $u_{\phi,\psi} \in X^{p,r}$. Thus the set of all solutions of the Schrödinger equation in $X^{p,r}$ equals the set of solutions $u_{\phi,\psi}$, $(\phi, \psi) \in Z$. As $T_-u_{\phi,\psi} = A_q(\phi, \psi)$, we obtain that the range of A_q equals to \mathcal{CD}_q .

Now, when the potential is equal to zero, the Dirichlet-to-Neumann operator $\Lambda_0 : u|_{\partial\Omega} \mapsto \partial_n u|_{\partial\Omega}$ is well-defined, $\Lambda_0 : W^{2-\frac{1}{r},r}(\partial\Omega) \rightarrow W^{1-\frac{1}{r},r}(\partial\Omega)$. The Cauchy data \mathcal{CD}_0 is the graph of the operator Λ_0 and is thus closed. Thus we see that the range of A_0 is a closed subspace, and therefore the operator $A_0 : Z/\ker(A_0) \rightarrow Z$ has zero kernel and closed range. Thus it is a semi-Fredholm operator. Now, consider the operator $A_q - A_0$. Using (3.9) we know that the operator

$$A_q - A_0 : Z/\ker(A_0) \rightarrow Z$$

is well-defined and compact. As compact perturbations of semi-Fredholm operators are also semi-Fredholm, we conclude that A_q is also semi-Fredholm.

It remains to show that $-A_q$ is a projection. This can be seen in a similar way to the smooth case. Indeed, if $(\phi, \psi) \in \text{ran}(A_q)$, $(\phi, \psi) = A_q(\tilde{\phi}, \tilde{\psi})$, we see that the solution $u_{\tilde{\phi}, \tilde{\psi}}$ has the trace $T_-u_{\tilde{\phi}, \tilde{\psi}} = (\phi, \psi)$. Hence Green's formula gives

$$u_{\tilde{\phi}, \tilde{\psi}} = -(-S_q \phi + K_q \psi) \quad \text{in } \Omega.$$

Taking the trace T_- from both sides, we obtain that $(\phi, \psi) = -A_q(\phi, \psi)$, i.e., $(-A_q)^2 = -A_q$. Thus, Proposition 3.3 is proven. \square

Now we can construct the boundary values of the exponentially growing solutions from the Cauchy data. Because we are given $\mathcal{CD}_q = \text{ran}(A_q)$ and we know $\ker(A_q) = \ker(A_0)$, we can construct the projection $-A_q$. Next, let $(\phi_\rho, \psi_\rho) = T_+e_\rho$ be the boundary values of the incoming plane wave. Consider the solution $v(x) = e^{\rho \cdot x}(1 + \psi(x, \rho)) = e^{\rho \cdot x} + u_0$, and let $(\phi, \psi) = T_+u_0$. Then $(\phi, \psi) \in \ker(A_q)$ and $u_0 = u_{\phi,\psi}$ in $\mathbb{R}^n \setminus \Omega$. Moreover, as v is a solution of the Schrödinger equation inside Ω , we have

$$(3.10) \quad T_+(e_\rho + u_{\phi,\psi}) = (\phi_\rho + \phi, \psi_\rho + \psi) + A_q(\phi, \psi) \in \text{ran}(A_q).$$

Applying the projection $I + A_q$ to (3.9) and using $A_q(\phi, \psi) = 0$, we see that

$$0 = (1 + A_q)(\phi_0 + \phi, \psi_0 + \psi) = (1 + A_q)(\phi_0, \psi_0) + (\phi, \psi).$$

Because A_q and (ϕ_0, ψ_0) are known, we find (ϕ, ψ) and the Cauchy data of $v(x)$ on $\partial\Omega$.

So far, we have constructed the Cauchy data of the solutions $v_{\rho_1}(x) = e^{\rho_1 \cdot x}(1 + \psi(x, \rho_1))$ for all sufficiently large ρ_1 . Thus, if we consider complex frequencies ρ_1 and ρ_2 satisfying $\rho_1 + \rho_2 = -i\xi$, with $\xi \in \mathbb{R}^n \setminus 0$, an application of Green's formula yields

$$\begin{aligned} \hat{q}(\xi) &= \lim_{|\rho_1| \rightarrow \infty} \int_{\Omega} q(x) e^{\rho_1 \cdot x} (1 + \psi_1(x, \rho_1)) \cdot e^{\rho_2 \cdot x} dx \\ &= \lim_{|\rho_1| \rightarrow \infty} \int_{\partial\Omega} (v_{\rho_1} \cdot \partial_n e^{\rho_2 \cdot x} - \partial_n v_{\rho_1} \cdot e^{\rho_2 \cdot x}) dx. \end{aligned}$$

This proves Theorem 3.1. \square

4 Nonuniqueness for Highly Singular Potentials

We next discuss how very strong singularities of the potential can cause non-uniqueness in a closely related inverse problem. Due to the strength of the singularities, the Schrödinger equation has to be interpreted in a weak sense. Let us consider the boundary value problem

$$(4.1) \quad (\Delta + q + E)u = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = f,$$

with the potential q having the form

$$(4.2) \quad q(x) = -\text{dist}(x, H)^\mu c_0(x)$$

near H , where dist is the Euclidean distance, H is a closed hypersurface bounding a region $\Omega_0 \Subset \Omega$, $\mu < -2$, and $c_0(x)$ is a smooth function, satisfying $c_0(x) > C_0 > 0$ in some neighborhood V of H .

Elements of $I^{-1-\mu}(H)$ satisfy the pointwise estimate $|q(x)| \leq C \text{dist}(x, H)^\mu$, but a q satisfying (4.2) is not even locally integrable and thus need not define a distribution. Hence, the solutions of (4.1) cannot be formulated in the usual sense of distributions. Instead, we define the solution of (4.1) (if it exists) to be the solution of the following convex minimization problem: Find u such that

$$(4.3) \quad G(u) = \inf G(v)$$

where $G = G_{q+E} : \{v \in H^1(\Omega) : v|_{\partial\Omega} = f\} \rightarrow \mathbb{R} \cup \{\infty\}$ is the convex functional

$$G_{q+E}(v) = \int_{\Omega} (|\nabla v(x)|^2 - (q(x) + E)|v(x)|^2) dx.$$

Here, since the function $-q(x)$ is bounded from below, we define $G(u) = \infty$ when $q|v|^2$ is not in $L^1(\Omega)$.

PROPOSITION 4.1 *The Cauchy data*

$$\mathcal{CD}_{q+E} = \left\{ \left(u|_{\partial\Omega}, \frac{\partial u}{\partial n} \Big|_{\partial\Omega} \right) : u \in H^1(\Omega), u \text{ is a minimizer of } G_{q+E} \right\}$$

does not depend on q in Ω_0 . In particular, if the solution of (4.3) is unique, u vanishes identically in Ω_0 .

Remark. We note that potentials having singularities similar to (4.2) as above have been used to produce counterexamples to strong unique continuation, e.g., potentials $q(x) = c/|x|^{2+\varepsilon}$ in [11]. Recently, counterexamples have been found for weak unique continuation for L^1 potentials [16], but here we need to construct potentials for which *all* solutions vanish inside H . Finally, we wish to emphasize that since the solutions of (4.1) considered here are not defined in the usual sense of distributions, but rather as solutions of a convex minimization problem, the solutions we construct do not give new counterexamples for the unique continuation problem.

PROOF: Obviously we can assume that $q(x) \leq 0$ everywhere. We start first with the case where $E = 0$ and $f \in C^\infty(\partial\Omega)$.

Because the potential q is not in the Kato class [6, p. 62], consider instead a decreasing sequence of smooth functions $q_n \in C^\infty(\Omega)$, $q_{n+1}(x) \leq q_n(x)$, for which $q_n(x) = q(x)$ when $d(x, H) > \frac{1}{n}$ and in some neighborhood V of H

$$(4.4) \quad q_n(x) \leq \max(-c_1 n^{-\mu}, q(x))$$

where $0 < c_1 < C_0$. Let G_n be the functionals defined as G with q replaced with q_n . The functionals G_n have unique minimizers u_n that satisfy in the classical sense

$$(4.5) \quad (\Delta + q_n)u_n = 0 \quad \text{in } \Omega, \quad u_n|_{\partial\Omega} = f.$$

Now, let $f \in C^\infty(\partial\Omega)$ be fixed. Let $F \in H^1(\Omega)$ be a function for which $F|_{\partial\Omega} = f$ and $F = 0$ in some neighborhood of H . By definition of the potentials q_n , for sufficiently large n_0 we have $G(F) = G_n(F) = G_{n_0}(F)$ for $n \geq n_0$. Thus for the minimizers u_n of G_n we have $G_n(u_n) \leq C = G_{n_0}(F)$. Next, by choosing a subsequence, we can assume that the sequences $\int |\nabla u_n(x)|^2 dx$ and $\int (-q_n(x))|u_n(x)|^2 dx$ are decreasing when $n \rightarrow \infty$. Next, let us denote by $C_1, C_2 \leq C$ the constants

$$C_1 = \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n(x)|^2 dx, \quad C_2 = \lim_{n \rightarrow \infty} \int_{\Omega} (-q_n(x))|u_n(x)|^2 dx = C_2.$$

Now, we see that u_n are uniformly bounded in $H^1(\Omega)$, and thus by choosing a subsequence we can assume that there is $\tilde{u} \in H^1(\Omega)$ such that $u_n \rightarrow \tilde{u}$ weakly in $H^1(\Omega)$. Moreover,

$$(4.6) \quad \int_{\Omega} |\nabla \tilde{u}(x)|^2 dx \leq C_1.$$

As compact operators map weakly converging sequences to strongly converging ones, we have the norm convergence $u_n \rightarrow \tilde{u}$ in $L^2(\Omega)$. Thus for $n_0 > 0$ we have

$$\begin{aligned} - \int_{\Omega} q_{n_0}(x) |\tilde{u}(x)|^2 dx &= \lim_{n \rightarrow \infty} \int_{\Omega} (-q_{n_0}(x)) |u_n(x)|^2 dx \\ &\leq \lim_{n \rightarrow \infty} \int_{\Omega} (-q_n(x)) |u_n(x)|^2 dx \leq C_2. \end{aligned}$$

Since this is valid for any n_0 , we have by the monotone convergence theorem

$$(4.7) \quad - \int_{\Omega} q(x) |\tilde{u}(x)|^2 dx \leq C_2.$$

Because $G \leq G_n$,

$$\inf G(u) \leq \lim_{n \rightarrow \infty} \min G_n(u) = C_1 + C_2.$$

Now, by (4.6) and (4.7) we have $G(\tilde{u}) \leq C_1 + C_2$, and thus \tilde{u} is a minimizer of G . Now, because $G = G_n + (G - G_n)$ where G_n is a strictly convex functional and $G - G_n$ is a convex functional, G is strictly convex. Thus the minimizer is unique. Hence we see that, for every f , the solution \tilde{u} of the minimization problem (4.3) exists, is unique, and is given as the L^2 limit of the functions u_n . We note that the above analysis was based on the fact that the minimization problems for the G_n epi-converge to the minimization problem for G [27].

Recalling that Ω_0 is the region bounded by H , consider functions u_n restricted to Ω_0 . Let $t \mapsto B_t$ be the Brownian motion in \mathbb{R}^n starting from x at time $t = 0$, i.e., $B_0 = x$. Because the q_n are strictly negative smooth functions, they are in the Kato class and the pair (Ω, q_n) is gaugeable (see [6, sect. 4.3 and theorem 4.19]). By [6, theorem 4.7], the solution u_n can be represented by the Feynman-Kac formula

$$u_n(x) = E(\exp\left(\int_0^\tau q_n(B_t) dt\right) f(B_\tau))$$

where $\tau = \tau_{\partial\Omega}$ is the first time when the process hits the boundary, i.e., $B_t \in \partial\Omega$. Here, we assume B_t is a version of Brownian motion for which all realizations are continuous curves (see [19] or [6, theorem 1.4]). If $x \in \Omega_0$, the realizations of Brownian motion have to hit H prior to hitting $\partial\Omega$. Denote the first hitting time for H by τ_H ; thus the first hitting point is B_{τ_H} , and $\tau_H < \tau_{\partial\Omega}$. (The stopping time τ_H is a measurable function in the probability space; see [6, prop. 1.15].)

Let us now denote by $p(\rho, \eta)$ the probability that the Brownian motion sent from the origin at time $t = 0$ leaves the origin-centered ball with radius ρ before time η . Because of the scale-invariance of Brownian motion, $p(s\rho, s^2\eta) = p(\rho, \eta)$ for $s > 0$. (Indeed, let us consider reparametrized Brownian motion $\tilde{B}_t = sB_{s^{-2}t}$. Because the probability densities of $(\tilde{B}_{t_1}, \dots, \tilde{B}_{t_m})$ coincide to those of $(B_{t_1}, \dots, B_{t_m})$, we see that \tilde{B}_t is Brownian motion, too.)

Let $A_{\rho,\eta} = \{|B_t - B_{\tau_H}| < \rho \text{ for } \tau_H \leq t < \tau_H + \eta\}$. This set is measurable in the probability space, and the probability of $A_{\rho,\eta}$ is $P(A_{\rho,\eta}) = 1 - p(\rho, \eta)$.

Let $m > 1$ and $\eta = \eta(m)$ be such that $p(1, \eta) \geq \frac{m-1}{m}$. Now q is nonpositive, and by (4.4) $q_n(x) < \max(-c_1 n^{-\mu}, q(x))$ in some neighborhood V of H . When s is so small that the s -neighborhood of H is in V , we have by (4.2) that

$$\begin{aligned} & \left| E \left(\exp \left(\int_0^\tau q(B_t) dt \right) f(B_\tau) \right) \right| \\ & \leq E \left(\exp \left(\int_{\tau_H}^{\tau_H + s^2 \eta} q(B_t) dt \right) \|f\|_{L^\infty} \right) \\ & \leq (1 - P(A_{s\rho, s^2 \eta(m)})) \|f\|_\infty \\ & \quad + P(A_{s\rho, s^2 \eta(m)}) \exp(-s^2 \eta(m) \min(C_0 s^\mu, c_1 n^{-\mu})) \|f\|_\infty. \end{aligned}$$

Thus, choosing $s = n^{2/\mu}$, we see that for sufficiently large n

$$(4.8) \quad \|u_n\|_{L^\infty(\Omega_0)} \leq \left(\frac{1}{m} + \frac{m-1}{m} \exp(-\eta(m) c_1 n^{2(2+\mu)/\mu}) \right) \|f\|_{L^\infty}.$$

Because $u_n \rightarrow \tilde{u}$ in $L^2(\Omega_0)$, $\|\tilde{u}\|_{L^2(\Omega_0)} \leq \frac{1}{m} \|f\|_{L^\infty} \text{vol}(\Omega_0)^{1/2}$ for any m . Thus we see that $\tilde{u} = 0$ in Ω_0 .

Next we consider the case when $E \in \mathbb{R}$ and $f \in H^{1/2}(\partial\Omega)$. First, let $H_r = \{x \in \Omega : \text{dist}(x, H) < r\}$, and let r be so small that $q(x) + E < 0$ for $x \in H_r$. If u is the solution of (4.1) in Ω , then its restriction $\tilde{u} = u|_{H_r}$ is the solution of the boundary value problem

$$(4.9) \quad (\Delta + q + E)\tilde{u} = 0 \quad \text{in } H_r, \quad \tilde{u}|_{\partial H_r} = \tilde{f},$$

where $\tilde{f} = u|_{\partial H_r} \in C^\infty(\partial H_r)$, that is, \tilde{u} is the solution of the minimization problem (4.3) in the domain H_r . Let q_n approximate q in H_r as above and \tilde{u}_n be the corresponding solutions of problem (4.9) with q replaced by q_n . As above, we see that problem (4.9) is uniquely solvable, $\tilde{u}_n \rightarrow \tilde{u}$ in $L^2(H_r)$, and that $\tilde{u}_n(x)$ can be represented by the Feynman-Kac formula. Let $x \in \Omega_0 \cap H_r$, $\tilde{\tau}$ be the first time when the Brownian motion sent from x at $t = 0$ hits ∂H_r , and $\tilde{A} = \{B_t \in \Omega_0 \cap H_r \text{ for } 0 \leq t < \tilde{\tau}\}$. Let $\tilde{f} = \tilde{f}_+ + \tilde{f}_-$, where \tilde{f}_+ vanishes on $\Omega_0 \cap \partial H_r$ and \tilde{f}_- vanishes on $(\Omega \setminus \Omega_0) \cap \partial H_r$. Then we see that

$$(4.10) \quad \begin{aligned} \tilde{u}_n(x) &= P(\tilde{A}) E \left(\exp \left(\int_0^{\tilde{\tau}} q_n(B_t) dt \right) \tilde{f}_-(B_{\tilde{\tau}}) \mid \tilde{A} \right) \\ &\quad + (1 - P(\tilde{A})) E \left(\exp \left(\int_0^{\tilde{\tau}} q_n(B_t) dt \right) \tilde{f}(B_{\tilde{\tau}}) \mid \tilde{A}^c \right), \end{aligned}$$

where $E(\cdot | \tilde{A})$ is conditional expectation with condition \tilde{A} , and \tilde{A}^c denotes the complement of \tilde{A} . Note that in the case of \tilde{A}^c , the process B_t hits H at least once. Analyzing how long Brownian motion is near H as above, we see that when $n \rightarrow \infty$,

the second term on the right-hand side of (4.10) goes to zero. Thus $\tilde{u}(x)$, for $x \in H_r \cap \Omega_0$, depends only on q in $H_r \cap \Omega_0$ and f_- . Similarly, we see that $\tilde{u}(x)$, $x \in H_r \setminus \overline{\Omega}_0$, depends only on q in $H_r \setminus \overline{\Omega}_0$ and f_+ . Moreover, analogously to (4.8) we see that

$$\lim_{n \rightarrow \infty} \|\tilde{u}_n|_H\|_{L^\infty(H)} = 0.$$

Choosing a subsequence, we can assume that $\tilde{u}_n \rightarrow \tilde{u}$ weakly in $H^1(H_r)$ and thus in $H^{3/4}(H_r)$. Hence, by taking the trace $H^{3/4}(H_r) \rightarrow L^2(H)$, we see that $\tilde{u}|_H = 0$.

In conclusion, for the boundary value problem (4.9) there are well-defined maps

$$\begin{aligned} T_+ : f_+ &\mapsto \tilde{u}|_{H_r \setminus \overline{\Omega}_0} \in \{v \in H^1(H_r \setminus \overline{\Omega}_0) : v|_H = 0\}, \\ T_- : f_- &\mapsto \tilde{u}|_{H_r \cap \Omega_0} \in \{v \in H^1(H_r \cap \Omega_0) : v|_H = 0\}, \end{aligned}$$

where T_+ depends only on q in $H_r \setminus \overline{\Omega}_0$ and T_- on q in $H_r \cap \Omega_0$.

In particular, on the boundaries $\partial H_r \cap \Omega_0$ and $\partial H_r \cap (\Omega \setminus \Omega_0)$, we have independent Dirichlet-to-Neumann maps

$$\Lambda_+ : f_+ \mapsto \partial_n \tilde{u}|_{\partial H_r \setminus \overline{\Omega}_0}, \quad \Lambda_- : f_- \mapsto \partial_n \tilde{u}|_{\partial H_r \cap \Omega_0},$$

where n is the exterior normal of H_r .

Next, if u is a solution of the boundary value problem (4.1), we denote $u_+ = u|_{\Omega \setminus \overline{\Omega}_0}$ and $u_- = u|_{\Omega_0}$. To motivate the next step, we observe that u_+ and u_- satisfy “independent” boundary value problems in $\Omega \setminus (H_r \cup \Omega_0)$,

$$(\Delta + q + E)u_+ = 0, \quad u_+|_{\partial H_r} = \tilde{f}, \quad \partial_n u_+|_{\partial H_r \setminus \overline{\Omega}_0} = \Lambda_+(u_+|_{\partial H_r \setminus \overline{\Omega}_0}),$$

and in $\Omega_0 \setminus H_r$

$$(\Delta + q + E)u_- = 0, \quad \partial_n u_-|_{\partial H_r \cap \Omega_0} = \Lambda_-(u_-|_{\partial H_r \cap \Omega_0}).$$

Now, considering the form of G and the fact that the solution u of boundary value problem (4.1) satisfies $u|_H = 0$, we see that $u_+ = u|_{\Omega \setminus \overline{\Omega}_0}$ is a minimizer of G in the set $\{v \in H^1(\Omega \setminus \overline{\Omega}_0) : v|_{\partial \Omega} = f, v|_H = 0\}$, and $u_- = u|_{\Omega_0}$ is a minimizer of G in the set $\{v \in H^1(\Omega_0) : v|_{\partial \Omega_0} = 0\}$.

Conversely, if $U = v_+$ in $\Omega \setminus \overline{\Omega}_0$ and $U = v_-$ in Ω_0 where v_+ and v_- are any minimizers of G in the sets $\{v \in H^1(\Omega \setminus \overline{\Omega}_0) : v|_{\partial \Omega} = f, v|_H = 0\}$ and $\{v \in H^1(\Omega_0) : v|_{\partial \Omega_0} = 0\}$, respectively, then U is a solution of (4.1).

In particular, we see that the Cauchy data of solutions u of (4.1) on $\partial \Omega$ are independent of $u|_{\Omega_0}$ and thus of q inside H . This finishes the proof of Proposition 4.1. As a concluding remark, we note that by using the Courant-Hilbert min-max principle, there always are values of E such that minimization problem for v_- has nonzero solutions; that is, there are eigenstates U that have vanishing Cauchy data on $\partial \Omega$. \square

Physically, this example has the following interpretation: In theory it is possible to construct a potential wall $q(x)$ such that no particles can “tunnel” through it, using an analogy with quantum mechanics. Thus exterior observers can make no

conclusions about the existence of objects or structures inside this wall. Moreover, inside H the solution can be in an eigenstate and its Cauchy data vanishes on the boundary of Ω . Thus, making another analogy with quantum mechanics, in this nest the Schrödinger cat could live happily ever after.

Acknowledgments. This work was supported in part by National Science Foundation grants DMS-9877101 and DMS-0070488 and Finnish Academy Project 172434. The third author was also supported by a John Simon Guggenheim Fellowship. The second and third author are grateful for the hospitality of the Mathematical Sciences Research Institute in Berkeley, California.

We would like to thank Steve McDowall for valuable discussions, and the referee for pointing out an error in the original version of this paper.

Bibliography

- [1] Beals, R.; Coifman, R. The spectral problem for the Davey-Stewartson and Ishimori hierarchies. *Nonlinear evolution equations: Integrability and spectral methods*, 15–23. Manchester University, Manchester, 1988.
- [2] Brown, R. M. Global uniqueness in the impedance-imaging problem for less regular conductivities. *SIAM J. Math. Anal.* **27** (1996), no. 4, 1049–1056.
- [3] Brown, R. M.; Uhlmann, G. A. Uniqueness in the inverse conductivity problem for nonsmooth conductivities in two dimensions. *Comm. Partial Differential Equations* **22** (1997), no. 5–6, 1009–1027.
- [4] Chanillo, S. A problem in electrical prospection and an n -dimensional Borg-Levinson theorem. *Proc. Amer. Math. Soc.* **108** (1990), no. 3, 761–767.
- [5] Chazarain, J.; Piriou, A. *Introduction to the theory of linear partial differential equations*. Studies in Mathematics and Its Applications, 14. North-Holland, Amsterdam–New York, 1982.
- [6] Chung, K. L.; Zhao, Z. X. *From Brownian motion to Schrödinger's equation*. Grundlehren der Mathematischen Wissenschaften, 312. Springer, Berlin, 1995.
- [7] Coifman, R. R.; McIntosh, A.; Meyer, Y. L'intégrale de Cauchy définit un opérateur borné sur L^2 pour les courbes lipschitziennes. *Ann. of Math. (2)* **116** (1982), no. 2, 361–387.
- [8] Colton, D.; Kress, R. *Inverse acoustic and electromagnetic scattering theory*. Second edition. Applied Mathematical Sciences, 93. Springer, Berlin, 1998.
- [9] Duistermaat, J. J.; Hörmander, L. Fourier integral operators. II. *Acta Math.* **128** (1972), no. 3–4, 183–269.
- [10] Faddeev, L. D. Growing solutions of the Schrödinger equation. *Dokl. Akad. Nauk SSR* **165** (1965), 514–517; translation in *Soviet Phys. Dokl.* **10** (1966), 1033.
- [11] Garofalo, N.; Lin, F.-H. Monotonicity properties of variational integrals, A_p weights and unique continuation. *Indiana Univ. Math. J.* **35** (1986), no. 2, 245–268.
- [12] Greenleaf, A.; Uhlmann, G. Estimates for singular Radon transforms and pseudodifferential operators with singular symbols. *J. Funct. Anal.* **89** (1990), no. 1, 202–232.
- [13] Greenleaf, A.; Uhlmann, G. Recovering singularities of a potential from singularities of scattering data. *Comm. Math. Phys.* **157** (1993), no. 3, 549–572.
- [14] Hörmander, L. *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis. II. Differential operators with constant coefficients. III. Pseudodifferential operators. IV. Fourier integral operators*. Grundlehren der Mathematischen Wissenschaften, 256, 257, 274, 275. Springer, Berlin, 1983, 1985.
- [15] Isakov, V. On uniqueness of recovery of a discontinuous conductivity coefficient. *Comm. Pure Appl. Math.* **41** (1988), no. 7, 865–877.

- [16] Kenig, C. E.; Nadirashvili, N. A counterexample in unique continuation. *Math. Res. Lett.* **7** (2000), no. 5-6, 625–630.
- [17] Knudsen, K.; Tamasan, A. Reconstruction of less regular conductivities in the plane. Preprint, 2001. Available online at <http://www.msri.org/publications/preprints/online/2001-035.html>.
- [18] Kohn, R. V.; Vogelius, M. Determining conductivity by boundary measurements. II. Interior results. *Comm. Pure Appl. Math.* **38** (1985), no. 5, 643–667.
- [19] Lévy, P. *Processus stochastiques et mouvement brownien*. Deuxième édition revue et augmentée. Gauthier-Villars, Paris, 1965.
- [20] Melrose, R. Marked lagrangians. Unpublished lecture notes, Massachusetts Institute of Technology.
- [21] Melrose, R.; Ritter, N. Interaction of nonlinear progressing waves for semilinear wave equations. *Ann. of Math. (2)* **121** (1985), no. 1, 187–213.
- [22] Nachman, A. I. Reconstructions from boundary measurements. *Ann. of Math. (2)* **128** (1988), no. 3, 531–576.
- [23] Nachman, A. I. Global uniqueness for a two-dimensional inverse boundary value problem. *Ann. of Math. (2)* **143** (1996), no. 1, 71–96.
- [24] Nachman, A.; Sylvester, J.; Uhlmann, G. An n -dimensional Borg-Levinson theorem. *Comm. Math. Phys.* **115** (1988), no. 4, 595–605.
- [25] Päivärinta, L.; Panchenko, A.; Uhlmann, G. Complex geometrical optics solutions for Lipschitz conductivities. *Rev. Mat. Iberoamericana*, in press.
- [26] Piriou, A. Calcul symbolique non linéaire pour une onde conormale simple. *Ann. Inst. Fourier (Grenoble)* **38** (1988), no. 4, 173–187.
- [27] Rockafellar, R. T.; Wets, R. J.-B. *Variational analysis*. Grundlehren der Mathematischen Wissenschaften, 317. Springer, Berlin, 1998.
- [28] Sylvester, J.; Uhlmann, G. A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math. (2)* **125** (1987), no. 1, 153–169.
- [29] Taylor, M. E. *Pseudodifferential operators and nonlinear PDE*. Progress in Mathematics, 100. Birkhäuser, Boston, 1991.

ALLAN GREENLEAF
 University of Rochester
 Department of Mathematics
 Rochester, NY 14627
 E-mail: allan@math.rochester.edu

MATTI LASSAS
 University of Helsinki
 Rolf Nevanlinna Institute
 P.O. Box 4 (Yliopistonkatu 5)
 FIN 00014 University of Helsinki
 FINLAND
 E-mail: Matti.Lassas@helsinki.fi

GUNTHER UHLMANN
 University of Washington
 Department of Mathematics
 Box 354350
 Seattle, WA 98195
 E-mail: gunther@math.washington.edu

Received December 2001.

Revised October 2002.