

# Home Exercises 6

28.10.2024

Write your name at the beginning of the file as “author:”.

1. Return to Moodle by **9.00am, Mon 28.10.** (to section “BEFORE”).
2. Watch the exercise session video available in Moodle by **10.00am, Mon 28.10.**
3. If you observe during the exercise session that your answers need some correction, return a corrected version to Moodle (to section “AFTER”) by **9.00 am, Mon 4.11.**

**Problem 1.** Read in file “systbp\_ldlc.txt”. (See Exercise set 1 for how to read it if you don’t remember.) Check whether there are any NAs in the data frame by command `anyNA( )`.

- (i) Plot `systbp` on x-axis and `ldlc` on y-axis and add their correlation value in the title of the plot.
- (ii) Fit a linear model of `ldlc ~ systbp`. Add the regression line to the existing plot. Show the `summary()` of the model fit. Is `systbp` a significant predictor of `ldlc` at significance level 0.05? Is it a useful predictor for any practical purposes, for example, could you use this model in a clinical setting to predict reliably `ldlc` from measured `systbp`? (Hint: Look at the plot and the R-squared value and think whether there is any useful predictive power here.)
- (iii) What is the 95%CI of the coefficient of `systbp` in this model?

**Problem 2.** Let’s consider the prostate cancer data set from Exercise set 4. Read it in by `y = read.table("prostate.txt", as.is = T, header = T)` and apply `head(y)`. Our variables are: log of cancer volume (`lcavol`), log prostate weight (`lweight`), age, log of benign prostatic hyperplasia amount (`lbph`), seminal vesicle invasion (`svi`), log of capsular penetration (`lcp`), Gleason score (`gleason`), percent of Gleason scores 4 or 5 (`pgg45`) and log(arithm) of PSA (`lpsa`).

- (i) Visualize the correlation matrix of the 9 variables mentioned above by `corrplot.mixed()` from package `corrplot` (See Lecture example 6.1.4.) You can pick the 9 variables by indexing columns with `2:10` rather than writing the column names explicitly.
- (ii) We want to predict `lcavol`. From `corrplot` we see that `lpsa` is a good candidate predictor as it is highly correlated with `lcavol`. Compute correlation with 95% confidence interval between `lcavol` and `lpsa`. For CI, use `r.con()` function from package `psych`.
- (iii) Fit a linear model `lcavol ~ lpsa`. Print out its `summary()`. How much variation in `lcavol` does the model explain?
- (iv) Plot values of `lpsa` on x-axis and `lcavol` on y-axis. Add the linear model fit to the same plot using `abline()` function.
- (v) Estimate visually from the plot what is an average `lcavol` for an individual whose `lpsa = 2.0`.
- (vi) Use the model coefficients from the linear model of (2.ii) to compute the exact linear model prediction for individual with `lpsa = 2.0`.

**Problem 3.** Continue with the prostate cancer data from Problem 2.

- (i) Fit a linear regression model for `lcavol` that has both variables `lpsa` and `lcp` included as predictors. How much variance does it explain?
- (ii) What is the formula by which the model from (3.i) turns the values of `lpsa` and `lcp` into the predicted value of `lcavol`?
- (iii) If an individual has values `lpsa=3` and `lcp=-1`, what is the predicted value for `lcavol`.

**Problem 4.** Continue with the prostate cancer data from Problem 2 & 3.

Split data into two parts based on the `lpsa` values of the individuals. Part I are individuals with `lpsa` at most the median `lpsa` in these data, and Part II are individuals with `lpsa` above the median `lpsa` in these data. Fit separate linear regression models in each Part of the data where you regress `lcavol` on `age` of the individuals. Do you notice differences how `age` predicts `lcavol` depending on whether `lpsa` values are low or high?