



Uniform Labelled Calculi for Conditional and Counterfactual Logics

Marianna Girlando^{1,2(✉)}, Sara Negri², and Giorgio Sbardolini³

¹ Aix Marseille Univ, Université de Toulon, CNRS, LIS, Marseille, France
marianna.girlando@univ-amu.fr

² University of Helsinki, Helsinki, Finland
sara.negri@helsinki.fi

³ The Ohio State University, Columbus, USA
sbardolini.1@osu.edu

Abstract. Lewis's counterfactual logics are a class of conditional logics that are defined as extensions of classical propositional logic with a two-place modal operator expressing conditionality. Labelled proof systems are proposed here that capture in a modular way Burgess's preferential conditional logic $\mathbb{P}CL$, Lewis's counterfactual logic \mathbb{V} , and their extensions. The calculi are based on preferential models, a uniform semantics for conditional logics introduced by Lewis. The calculi are analytic, and their completeness is proved by means of countermodel construction. Due to termination in root-first proof search, the calculi also provide a decision procedure for the logics.

Keywords: Conditional logics · Counterfactual logics · Proof theory · Preferential models · Labelled calculi

1 Introduction

In Stalnaker's and Lewis's approach, conditional logics are defined as extensions of classical propositional logic by means of a two-place modal operator, the conditional, here denoted as $>$. This intensional operator is intended to express a more fine-grained notion of conditionality than material implication.

Lewis introduced counterfactual conditional logics to extend formal reasoning to counterfactual sentences, i.e., statements of the form *If Trump hadn't won the elections, Clinton would have been president*. Other than counterfactual logics (system \mathbb{V} and its extensions) conditional logics include a weaker family of systems: in this paper, we consider *preferential conditional logic* $\mathbb{P}CL$ and all its extensions. These latter systems have received attention in artificial intelligence since the conditional operator can be interpreted as expressing non monotonic

This work was partially supported by the Academy of Finland research project no. 1308664 and by the project TICAMORE ANR-16-CE91-0002-01.

© Springer-Verlag GmbH Germany, part of Springer Nature 2019
R. Iemhoff et al. (Eds.): WoLLIC 2019, LNCS 11541, pp. 1–16, 2019.
https://doi.org/10.1007/978-3-662-59533-6_16

inferences, i.e., sentences of the form *Normally, cats are afraid of dogs*. In particular, the fragment of $\mathbb{P}\text{CL}$ without nesting of the conditional operator is equivalent to system \mathbf{P} of [14]. There are other applications of conditional logics in the fields of knowledge base update [13], causality [8] and, in an epistemic setting, belief revision [4, 12].

The semantics of conditional logics is defined in terms of possible world structures in which, intuitively, a formula $A > B$ is true at world x if B is true in the set of worlds at which A is true that are more similar, in a sense to be formalized, to x . Preferential models were proposed by Lewis and studied, among others, by Burgess, who proved completeness of $\mathbb{P}\text{CL}$ with respect to these models [5]. Halpern and Friedman extended the proof to extensions of $\mathbb{P}\text{CL}$ [7]. On a formal level, these models explicitly employ the notion of comparative similarity among worlds: they are defined by adding to a set of possible worlds W a family of subsets W_x for each $x \in W$, representing the worlds accessible from x , and a binary relation \leq_x , expressing similarity among worlds. Thus, $y \leq_x z$ means *world y is at least as similar as z to world x* .

In this article, we define a family of modular labelled calculi $\mathbf{G3P}^*$ for conditional logic $\mathbb{P}\text{CL}$ and all its extensions, including counterfactual logics, i.e., \mathbb{V} and its extensions. The calculi are based on preferential semantics: following the well-established methodology proposed by the second author, the calculi import into the sequent calculus the semantic elements of preferential models by means of syntactic elements (labels and relational symbols).

In [20], the second and third author presented a labelled calculus based on ternary relations for a system of Lewis's conditional logic $\mathbb{V}\text{C}$ (\mathbb{V} to which the condition of Centering is added). The present article stems from a comment in Weiss's thesis [22]: the author observes, correctly, that Negri and Sbardolini's proof system is actually adequate to capture the stronger system $\mathbb{V}\text{CU}$ ($\mathbb{V}\text{C} + \text{Uniformity}$). This led us to an analysis of labelled calculi based on preferential models. It turns out that it is possible to define modular proof systems on the basis of these natural classes of models.

The article is organised as follows. In Sect. 2, conditional logics and preferential models are introduced. Section 3 presents the rules of the calculi $\mathbf{G3P}^*$ and Sect. 4 their structural properties. In Sect. 5, we define a proof search strategy that ensures termination in root-first proof search for the systems without the semantic conditions of Uniformity and Absoluteness. This allows to prove completeness of the calculi by extracting a countermodel from failed proof search. The conclusion (Sect. 6) gives a comparison of the calculi presented in this article with other proof systems for conditional logics found in the literature.

2 Conditional Logics and Preferential Models

The language of conditional logics is defined by means of the following grammar, for p propositional variable, $A, B \in \mathcal{L}_{\text{cond}}$, and $>$ the conditional operator:

$$\mathcal{L}_{\text{cond}} = p \mid \perp \mid A \wedge B \mid A \vee B \mid A \rightarrow B \mid A > B$$

An axiomatization of $\mathbb{P}CL$ is defined adding to the axioms and inference rules of classical propositional logic the following:

$$\begin{array}{ll}
\text{RCEA} & \frac{A \leftrightarrow B}{(A > C) \leftrightarrow (B > C)} & \text{RCK} & \frac{A \rightarrow B}{(C > A) \rightarrow (C > B)} \\
\text{R-And} & (A > B) \wedge (A > C) \rightarrow (A > (B \wedge C)) & \text{ID} & A > A \\
\text{CM} & (A > B) \wedge (A > C) \rightarrow ((A \wedge B) > C) & \text{RT} & (A > B) \wedge ((A \wedge B) > C) \rightarrow (A > C) \\
\text{OR} & (A > C) \wedge (B > C) \rightarrow ((A \vee B) > C)
\end{array}$$

An axiomatization of Lewis's logic \mathbb{V} is obtained by adding axiom CV to $\mathbb{P}CL$. Extensions of both $\mathbb{P}CL$ and \mathbb{V} are defined by adding the axioms for *normality*, *total reflexivity*, *weak centering*, *centering*, *uniformity* and *absoluteness* (denoted below by the corresponding initials). The family of systems is represented in Fig. 1.

$$\begin{array}{ll}
\text{CV} & ((A > C) \wedge \neg(A > \neg B)) \rightarrow ((A \wedge B) > C) \\
\text{N} & \neg(\top > \perp) & \text{T} & A \rightarrow \neg(A > \perp) \\
\text{W} & (A > B) \rightarrow (A \rightarrow B) & \text{C} & (A \wedge B) \rightarrow (A > B) \\
\text{U}_1 & (\neg A > \perp) \rightarrow \neg(\neg A > \perp) > \perp & \text{U}_2 & \neg(A > \perp) \rightarrow ((A > \perp) > \perp) \\
\text{A}_1 & (A > B) \rightarrow (C > (A > B)) & \text{A}_2 & \neg(A > B) \rightarrow (C > \neg(A > B))
\end{array}$$




Fig. 1. The conditional logics cube

We use preferential models as the semantics for the conditional logics cube. Preferential models were studied, among others, by Burgess, who proved the adequacy of this class of models with respect to preferential logic [5].

Definition 1. A preferential model $\langle W, \{W_x\}_{x \in W}, \{\leq_x\}_{x \in W}, [\] \rangle$ consists of a non-empty set of worlds, W , and for every world $x \in W$, a set W_x of worlds accessible from it, and a binary relation \leq_x over W . The propositional evaluation

$\llbracket \cdot \rrbracket : \text{Atm} \rightarrow \mathcal{P}(W)$ selects the worlds at which an atomic formula is true. The relation \leq_x satisfies the following properties:

- Reflexivity, for all $w \in W$, $w \leq_x w$, and
- Transitivity, for all $w, y, z \in W$, if $w \leq_x y$ and $y \leq_x z$ then $w \leq_x z$.

The truth condition for the conditional operator within preferential models is:

$$x \Vdash A > B \equiv \text{for all } z \in W_x, \text{ if } z \Vdash A, \text{ then there exists } y \in W_x \text{ such that } y \leq_x z, y \Vdash A, \text{ and for all } k \in W_x, \text{ if } k \leq_x y \text{ then } k \Vdash A \rightarrow B.$$

Extensions of preferential models are specified by adding conditions on the relation \leq_x . These models are adequate for the logics in the conditional cube [7].

Definition 2. *Extensions of preferential models are defined as follows:*

- Normality: For all $x \in W$, W_x is non-empty;
- Total reflexivity: For all $x \in W$ it holds that $x \in W_x$;
- Weak centering: For all $x \in W$, for all $y \in W_x$, it holds that $x \leq_x y$;
- Centering: For all $x \in W$, for all $y \in W_x$, it holds that $x \leq_x y$ and if there is $w \in W_x$ such that for all $y \in W_x$, $w \leq_x y$, then $w = x$;
- Uniformity: For all $x \in W$, for all $y \in W_x$ it holds that $W_y = W_x$;
- Absoluteness: Uniformity plus for all $w_1, w_2 \in W_x$, $w_1 \leq_x w_2$ iff $w_1 \leq_y w_2$;
- Nesting: For all $x \in W$, for all $w_1, w_2 \in W_x$, either $w_1 \leq_x w_2$ or $w_2 \leq_x w_1$.

Some of the above conditions are incremental: total reflexivity implies normality, weak centering implies total reflexivity, centering implies weak centering and absoluteness implies uniformity.

3 Labelled Proof Systems

In this section we shall define a family of modular calculi for the conditional cube. To this aim, we enrich our language with a sets of labels x, y, z, \dots denoting worlds in preferential models. Furthermore, we allow the following expressions to occur in sequents: labelled formulas $x : A$, denoting $x \Vdash A$, and relational atoms $y \in W_x$, $y \leq_x z$ and $x = y$, having the same meaning as their semantic counterparts. Following [20], we introduce an indexed modal operator and reformulate the truth condition of $A > B$ in terms of this operator:

$$w \Vdash \Box_x A \equiv \text{for all } k \in W_x, \text{ if } k \leq_x w \text{ then } k \Vdash A$$

$$(*) x \Vdash A > B \equiv \text{for all } z \in W_x, \text{ if } z \Vdash A, \text{ then there exists } y \in W_x \text{ such that } y \leq_x z, y \Vdash A, \text{ and } y \Vdash \Box_x(A \rightarrow B).$$

Following [19], we introduce an indexed conditional operator to treat the second disjunct of the truth condition of the conditional operator:

$$C_x^z(A, B) \equiv \text{there exists } y \in W_x \text{ such that } y \leq_x z, y \Vdash A \text{ and } y \Vdash \Box_x(A \rightarrow B)$$

Thus, the truth condition for the conditional operator can be stated as follows:

$$(**) x \Vdash A > B \equiv \text{for all } z \in W_x, \text{ if } z \Vdash A, \text{ then } C_x^z(A, B).$$

Observe that the extension of the language is only at the level of the labelled rules and produces, in the course of proof search, only formulas of a certain specific form. Formulas containing the new operators never occur as proper subformulas of other formulas and an indexed modality can have only an implication in its scope.

The rules of the labelled proof systems are defined by analysing the truth conditions of the above operators (Fig. 2). Rules Ref and Tr express reflexivity and transitivity of \leq_x ; rule Ref₌ and Repl express reflexivity of equality and the property of replacement of equals. We call **G3P** the calculus for \mathbb{PCL} ; calculi for extensions are defined in a modular way by adding to \mathbb{PCL} the rules corresponding to the semantic properties of \leq_x . We denote by **G3P*** the whole family of calculi. The condition of freshness of a variable y in a rule is indicated by $(y!)$.

In order to prove soundness of the rules of the calculus we have to provide a definition of realization in a preferential model. The definition uses the operators of the extended language, and to guarantee its non-circularity we need to define a notion of weight of formulas:

Definition 3. *Given a labelled formula F of the form $x : A$, let the pure part of F be defined as $p(x : A) = A$, and the labelled part as $l(x : A) = l(x : \Box_k A) = x$. The weight of a labelled formula is an ordered pair $\langle w(p(F)), w(l(F)) \rangle$ where*

- for x world label, $w(x) = 0$
- $w(p) = w(\perp) = 1$; $w(A \circ B) = w(A) + w(B) + 1$, for \circ conjunction, disjunction or implication; $w(x : \Box_k A) = w(A) + 1$; $w(C_x^z(A, B)) = w(A) + w(B) + 3$; $w(A > B) = w(A) + w(B) + 4$.

Definition 4 (Realization). *Given a model $\mathcal{M} = \langle W, \{W_x\}_{x \in W}, \{\leq_x\}_{x \in W}, \llbracket \rrbracket \rangle$, and a set P of world labels, a P -realization over \mathcal{M} is a function $\rho : P \rightarrow W$ that assigns to each world label $x \in P$ an element $\rho(x) \in W$. Satisfiability of a formula $F \in \mathcal{L}_{\text{cond}}$ is defined by cases as follows: $\mathcal{M} \models_\rho y \in W_x$ if $\rho(y) \in W_{\rho(x)}$; $\mathcal{M} \models_\rho y \leq_x z$ if $\rho(y) \leq_{\rho(x)} \rho(z)$; $\mathcal{M} \models_\rho x : p$ if $\rho(x) \in \llbracket p \rrbracket$, for p atomic;¹ $\mathcal{M} \models_\rho w : \Box_x A$ if for all $y \in W_{\rho(x)}$, if $y \leq_{\rho(x)} \rho(w)$, then $y \Vdash A$; $\mathcal{M} \models_\rho C_x^z(A, B)$ if there exists $y \in W_{\rho(x)}$ such that $y \leq_x \rho(z)$, $y \Vdash A$ and $y \not\vdash \Box_x(A \rightarrow B)$; $\mathcal{M} \models_\rho x : A > B$ if for all $k \in W_{\rho(x)}$, if $k \Vdash A$, then $k \not\vdash C_x^k(A, B)$. A sequent $\Gamma \Rightarrow \Delta$ is valid in \mathcal{M} under the ρ realization iff whenever $\mathcal{M} \models_\rho F$ for all $F \in \Gamma$, then $\mathcal{M} \models_\rho G$ for some $G \in \Delta$. A sequent is valid in a class of preferential models if it is valid under any realization for any model of that class.*

The above definition immediately yields:

¹ The definition can be extended to the propositional formulas of the language in the standard way [17].

Initial sequents		
$x : p, \Gamma \Rightarrow \Delta, x : p$	$x : \perp, \Gamma \Rightarrow \Delta$	
Propositional rules (standard)		
Conditional rules		
$\frac{k \in W_x, k \leq_x w, w : \Box_x A, k : A, \Gamma \Rightarrow \Delta}{k \in W_x, k \leq_x w, w : \Box_x A, \Gamma \Rightarrow \Delta} \text{L}\Box_x$	$\frac{k \in W_x, k \leq_x w, \Gamma \Rightarrow \Delta, k : A}{\Gamma \Rightarrow \Delta, w : \Box_x A} \text{R}\Box_x(k!)$	
$\frac{y \in W_x, y \leq_x z, y : A, y : \Box_x(A \rightarrow B), \Gamma \Rightarrow \Delta}{C_x^z(A, B), \Gamma \Rightarrow \Delta} \text{LC}(y!)$		
$\frac{y \leq_x z, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta, C_x^z(A, B), y : A \quad y \leq_x z, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta, C_x^z(A, B), y : \Box_x(A \rightarrow B)}{y \leq_x z, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta, C_x^z(A, B)} \text{RC}$		
$\frac{z \in W_x, x : A > B, \Gamma \Rightarrow \Delta, z : A \quad z \in W_x, x : A > B, C_x^z(A, B), \Gamma \Rightarrow \Delta}{z \in W_x, x : A > B, \Gamma \Rightarrow \Delta} \text{L} >$		
$\frac{z \in W_x, z : A, \Gamma \Rightarrow \Delta, C_x^z(A, B)}{\Gamma \Rightarrow \Delta, x : A > B} \text{R} >(z!)$		
Relational rules		
$\frac{w \leq_x w, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta} \text{Ref}$	$\frac{w \leq_x z, w \leq_x y, y \leq_x z, \Gamma \Rightarrow \Delta}{w \leq_x y, y \leq_x z, \Gamma \Rightarrow \Delta} \text{Tr}$	
$\frac{x = x, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta} \text{Ref}=_$	$\frac{x = y, \text{At}(x), \text{At}(y), \Gamma \Rightarrow \Delta}{x = y, \text{At}(y), \Gamma \Rightarrow \Delta} \text{Repl}$	
Rules for extensions		
$\frac{y \in W_x, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta} \text{N}(y!)$	$\frac{x \in W_x, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta} \text{T}$	$\frac{x \leq_x y, y \in W_x, \Gamma \Rightarrow \Delta}{y \in W_x, \Gamma \Rightarrow \Delta} \text{W}$
$\frac{x = y, y \leq_x x, y \in W_x, \Gamma \Rightarrow \Delta}{y \leq_x x, y \in W_x, \Gamma \Rightarrow \Delta} \text{C}$		
$\frac{z \in W_x, y \in W_x, z \in W_y, \Gamma \Rightarrow \Delta}{y \in W_x, z \in W_y, \Gamma \Rightarrow \Delta} \text{U1}$		$\frac{z \in W_y, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta}{y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta} \text{U2}$
$\frac{y \leq_x z, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta \quad z \leq_x y, y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta}{y \in W_x, z \in W_x, \Gamma \Rightarrow \Delta} \text{Nes}$		$\frac{y \leq_k z, y \leq_x z, \Gamma \Rightarrow \Delta}{y \leq_x z, \Gamma \Rightarrow \Delta} \text{A}$
<i>At(y) denotes any atoms of the form $y : p, y \in W_x, x \in W_y, y \leq_x z, x \leq_y z$.</i>		

G3P = Initial sequents, propositional rules, conditional rules, relational rules

G3P^N = **G3P** + N; **G3P^T** = **G3P^N** + T; **G3P^W** = **G3P^T** + W; **G3P^C** = **G3P^W** + C

G3P^U = **G3P** + U1 + U2; **G3P^{NU/TU/WU/CU}** = **G3P^{N/T/W/C}** + U1 + U2

G3P^A = **G3P** + A; **G3P^{NA/TA/WA/CA}** = **G3P^{N/T/W/C}** + A

G3P^V = **G3P** + Nes; **G3P^{VN/VT/VW/VCU}** = **G3P^{N/T/W/C}** + Nes

G3P^{VNU/VTU/VWU/VCU} = **G3P^{NU/TU/WU/CU}** + Nes;

G3P^{VNA/VTa/VWA/VCA} = **G3P^{NA/TA/WA/CA}** + Nes

Fig. 2. Rules of **G3P***

Theorem 1 (Soundness). *If a sequent is derivable in **G3P***, then it is valid in the corresponding class of preferential models.*

Remark 1. The sequent calculi of Fig. 2 are fully modular. However, by dropping the requirement of modularity, it is possible to define simpler versions of the calculi for some subfamilies of the logics. In systems with uniformity it holds that for all $x \in W$ and $y \in W_x$, $W_x = W_y$. Thus, we can avoid specifying the relational atoms $y \in W_x$ in the rules (with this reformulation, the rules of uniformity would become superfluous). Similarly, in logics with absoluteness, uniformity holds, and moreover $w_1 \leq_x w_2$ iff $w_2 \leq_y w_1$. Thus, we can avoid specifying the subscript x in relational atoms $y \leq_x z$, and the rule of absoluteness becomes superfluous. Finally, in the presence of nesting the truth condition for the conditional operator can be stated in a simpler way:

$x \Vdash A > B \equiv$ if there exists $z \in W_x$ such that $z \Vdash A$, then there exists $y \in W_x$ such that $y \Vdash A$ and $y \Vdash \Box_x(A \rightarrow B)$.

Rules based on this truth condition, in addition to the simplification explained for uniformity, i.e., no relational atoms $y \in W_x$ and no rules U1, U2, yield the calculus proposed in [20], a proof system sound and complete with respect to the conditional logic $\forall\text{CU}$.

4 Structural Properties

The *height* of a derivation is the number of nodes of the longest derivation branch, minus one. We recall that a rule is *height-preserving admissible* if whenever its premiss is derivable, the conclusion is also derivable with no greater derivation height. A rule is *height-preserving invertible* if whenever its conclusion is derivable, the premisses are derivable with no greater derivation height. Derivability with height bounded by n is denoted by \vdash_n .

In order to prove admissibility of the structural rules we need a notion of label substitution given by, for instance, $x : A > B[y/x] \equiv y : A > B$ and $w : \Box_x A[y/x] \equiv w : \Box_y A$, extended component-wise to sequents, and a property of height-preserving substitution: If $\vdash_n \Gamma \Rightarrow \Delta$, then $\vdash_n \Gamma[y/x] \Rightarrow \Delta[y/x]$. Admissibility of generalized initial sequents (i.e., sequents of the form $x : A, \Gamma \Rightarrow \Delta, x : A$ in which A is not necessarily atomic) is shown by induction on the weight of A . We omit the routine proofs, the details of which are similar to those in [20].

The *structural rules* of weakening, contraction, and cut of **G3P*** are the following:

$$\frac{\Gamma \Rightarrow \Delta}{\mathcal{F}, \Gamma \Rightarrow \Delta} \text{Wk}_L \quad \frac{\Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta, \mathcal{F}} \text{Wk}_R \quad \frac{\mathcal{F}, \mathcal{F}, \Gamma \Rightarrow \Delta}{\mathcal{F}, \Gamma \Rightarrow \Delta} \text{Ctr}_L \quad \frac{\Gamma \Rightarrow \Delta, \mathcal{F}, \mathcal{F}}{\Gamma \Rightarrow \Delta, \mathcal{F}} \text{Ctr}_R$$

$$\frac{\Gamma \Rightarrow \Delta, \mathcal{F} \quad \mathcal{F}, \Gamma' \Rightarrow \Delta'}{\Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \text{cut}$$

where \mathcal{F} is a relational atom, a labelled formula, or a formula of the form $C_x^z(A, B)$. Observe that for Wk_R , Ctr_R , and cut we can without loss of generality omit the case of relational formulas since they never occur in the right-hand side of sequents. The calculi **G3P*** have the following structural properties:

Theorem 2.

- i. All the rules are height-preserving invertible.
- ii. The rules of weakening and contraction are height-preserving admissible.
- iii. The rule of cut is admissible.

Proof.

i. By induction on the height of the derivation. Invertibility of relational rules, rules for extensions, $\text{L}\Box_x$, RC and $\text{L}>$ immediately follows from admissibility of weakening. Invertibility of the propositional rules and of $\text{R}\Box_x$ is proved as in [17]; invertibility of LC is similar to that of the corresponding rule in [19].

ii. By induction on the height n of the derivation. If $n = 0$, the premiss of the contraction rule is an initial sequent, and so is its conclusion. If $n > 0$, we look at the last rule (R) applied. If F is not principal in the rule, it suffices to apply the inductive hypothesis to the premiss of (R) , and then (R) . If F is the principal formula of R , or was introduced by R , we distinguish two subcases. If (R) is a rule in which the principal formula appears also in the premiss apply the hypothesis to the premiss, and then the rule. If (R) is a rule in which the active formulas are subformulas of the principal formula, apply invertibility to the premiss(es) of the rule, then the inductive hypothesis, and (R) .

iii. By primary induction on the weight of the cut formula, and secondary induction on the sum of heights of the derivations of the premisses of cut. As usual, we proceed with a case distinction according to the last rule applied. If at least one of the premisses is an initial sequent, the conclusion of cut is also a sequent, or can be obtained by easy rule permutations. Similarly, if the cut formula is not principal in the last rule R applied to one of the premiss of cut, the conclusion of cut can be obtained by permuting the cut upwards on the premiss of R , and then applying R again. Finally, if the cut formula is principal in both rules applied to the premisses of cut, some more complex permutations are needed. Propositional cases can be found in [20]. We show only the case in which $R >$ and $L >$ are the rules applied to the left and right premiss of cut respectively. Consider a derivation ending with

$$\frac{\frac{y \in W_x, y : A, \Gamma \Rightarrow \Delta, C_x^y(A, B)}{\Gamma \Rightarrow \Delta, x : A > B} \text{ R} > \quad \frac{z \in W_x, x : A > B, \Gamma' \Rightarrow \Delta', z : A \quad z \in W_x, x : A > B, C_x^y(A, B), \Gamma' \Rightarrow \Delta'}{z \in W_x, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \text{ L} >}{z \in W_x, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'} \text{ cut}$$

Let $\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3$ be the derivations ending with the topsequents above. The cut is transformed into four cuts of reduced rank as follows. First we have two cuts, to topmost of reduced height, the second of reduced weight, where $\mathcal{D}_1[z/y]$ denotes the derivation resulting from \mathcal{D}_1 by application of an height-preserving substitution:

$$\frac{\frac{\Gamma \Rightarrow \Delta, x : A > B \quad z \in W_x, x : A > B, \Gamma' \Rightarrow \Delta', z : A}{z \in W_x, \Gamma, \Gamma' \Rightarrow \Delta, \Delta', z : A} \text{ cut} \quad \mathcal{D}_1[z/y]}{z \in W_x^2, \Gamma^2, \Gamma' \Rightarrow \Delta^2, \Delta', C_x^z(A, B)} \text{ cut}$$

Second we have the cut of reduced height

$$\frac{\Gamma \Rightarrow \Delta, x : A > B \quad \mathcal{D}_3}{z \in W_x, \Gamma, \Gamma', C_x^z(A, B) \Rightarrow \Delta, \Delta'} \text{ cut}$$

Finally, by cut their conclusions through a fourth cut of reduced weight and obtain the sequent $z \in W_x^3, \Gamma^3, \Gamma'^2 \Rightarrow \Delta^3, \Delta'^2$. Admissible weakening steps give the conclusion of the original cut.

The case of principal cut formula of the form $C_x^z(A, B)$ is reduced in a similar way through four cuts, the uppermost of reduces height, and the lowermost of reduces weight. For principal formula of the form of an indexed modality, the conversion is the standard one for the necessity modality of labelled calculi.

Thanks to admissibility of cut, it is possible to prove the following:

Theorem 3 (Completeness). *If a formula A is valid in preferential models and extensions, then sequent $\Rightarrow x : A$, for an arbitrary label x , is derivable in the corresponding $\mathbf{G3P}^*$ calculus.*

Proof. By using the known completeness result for extensions of PCL w.r.t. preferential models and showing that the inference rules of PCL are admissible in $\mathbf{G3P}$, and that the axioms of PCL and its extensions are derivable in the corresponding proof system of $\mathbf{G3P}^*$. The proof for PCL is similar to the proof in [20]. By way of example, we show the derivation of Axiom U_1 in $\mathbf{G3P}^{\text{VU}}$, omitting the derivable left premiss of $L >$.

$$\begin{array}{c}
 \frac{y : \perp \dots \Rightarrow \dots \quad \dots \quad y : \neg A \Rightarrow y : \neg A \dots}{y \leq_x y, y \leq_x w, y \in W_x, \dots, y : \neg A, y : \neg A \rightarrow \perp \Rightarrow \dots} L_{\rightarrow} \\
 \frac{k \leq_x k, k \leq_x w, k \in W_x, w \in W_x, w \in W_z, z \in W_x, w : \neg A, w : \neg A, k : \neg A, k : \Box_x(\neg A \rightarrow \perp), x : (\neg A > \perp) \Rightarrow \dots}{k \leq_x w, k \in W_x, w \in W_x, w \in W_z, z \in W_x, w : \neg A, w : \neg A, k : \neg A, k : \Box_x(\neg A \rightarrow \perp), x : (\neg A > \perp) \Rightarrow \dots} \text{Ref} \\
 \frac{\dots}{w \in W_x, w \in W_z, z \in W_x, w : \neg A, w : \neg A, C_x^w(\neg A, \perp), x : (\neg A > \perp) \Rightarrow \dots} LC \\
 \frac{\dots}{w \in W_x, w \in W_z, z \in W_x, w : \neg A, x : (\neg A > \perp) \Rightarrow \dots} L > \\
 \frac{\dots}{w \in W_z, z \in W_x, w : \neg A, x : (\neg A > \perp) \Rightarrow C_x^z(\neg(\neg A > \perp), \perp), C_z^w(\neg A, \perp)} U1 \\
 \frac{\dots}{z \in W_x, x : (\neg A > \perp) \Rightarrow z : \neg A > \perp, C_x^z(\neg(\neg A > \perp), \perp)} R > \\
 \frac{\dots}{z \in W_x, z : \neg(\neg A > \perp), x : (\neg A > \perp) \Rightarrow C_x^z(\neg(\neg A > \perp), \perp)} R_{\neg} \\
 \frac{\dots}{x : (\neg A > \perp) \Rightarrow x : \neg(\neg A > \perp) > \perp} R > \\
 \frac{\dots}{\Rightarrow x : (\neg A > \perp) \rightarrow \neg(\neg A > \perp) > \perp} R_{\rightarrow}
 \end{array}$$

By Theorems 1 and 3, and the known completeness results for PCL and its extensions with respect to preferential models, we have:

Corollary 1. *Formula A is provable in any of the systems of the conditional logics cube if and only if $\Rightarrow x : A$ is derivable in the corresponding labelled system.*

5 Termination and Completeness

In this section, we shall give an alternative direct proof of completeness for the calculi $\mathbf{G3P}$, $\mathbf{G3P}^{\text{N}}$, $\mathbf{G3P}^{\text{T}}$, $\mathbf{G3P}^{\text{W}}$, $\mathbf{G3P}^{\text{C}}$ and $\mathbf{G3P}^{\text{V}}$, $\mathbf{G3P}^{\text{VN}}$, $\mathbf{G3P}^{\text{VT}}$, $\mathbf{G3P}^{\text{VW}}$, $\mathbf{G3P}^{\text{VC}}$ (from now on $\mathbf{G3P}^{\text{V/N/T/W/C}}$), i.e., the systems without uniformity and absoluteness.² The proof proceeds by showing how to construct a countermodel from failed proof search. We first need to prove that root-first proof search, which in general is not terminating because of loops, terminates under the adoption of a suitable strategy.

Example 1. Loop generated by repeated applications of rule $L >$ and LC (only the right premisses of $L >$ are shown).

² The proofs of termination and completeness for systems with Uniformity and Absoluteness can be given adopting the reformulation of the calculi from Remark 1. The proofs for the current versions of the calculi would be unnecessarily complex.

$$\begin{array}{c}
\vdots \\
y \leq_x z, y \in W_x, z \leq_x z, z \in W_x, y : A, y : \Box_x(A \rightarrow B), x : A > B, x : C > D, C_x^z(C, D), C_x^y(C, D) \Rightarrow \\
\frac{y \leq_x z, y \in W_x, z \leq_x z, z \in W_x, y : A, y : \Box_x(A \rightarrow B), x : A > B, x : C > D, C_x^z(C, D) \Rightarrow}{z \leq_x z, z \in W_x, x : A > B, x : C > D, C_x^z(A, B), C_x^z(C, D) \Rightarrow} \text{LC} \\
\frac{z \leq_x z, z \in W_x, x : A > B, x : C > D, C_x^z(A, B), C_x^z(C, D) \Rightarrow}{z \in W_x, x : A > B, x : C > D, C_x^z(A, B), C_x^z(C, D) \Rightarrow} \text{Ref} \\
\frac{z \in W_x, x : A > B, x : C > D, C_x^z(A, B) \Rightarrow}{z \in W_x, x : A > B, x : C > D \Rightarrow} \text{L} >
\end{array}$$

To ensure termination we introduce the notion of *saturated sequent*, a sequent to which all the rules have been applied in a non-redundant way. We then specify a *proof search strategy*, blocking the application of the rules to a saturated sequent.

Definition 5. Given a $\mathbf{G3P}^{\mathbf{V}/\mathbf{N}/\mathbf{T}/\mathbf{W}/\mathbf{C}}$ derivation, let $\mathcal{B} = S_0, S_1, \dots$ be a derivation branch, with S_i sequent $\Gamma_k \Rightarrow \Delta_k$ for $k > 0$, and S_0 sequent $\Rightarrow x_0 : A_0$. Let $\downarrow \Gamma_k / \downarrow \Delta_k$ be the union of the antecedents/succedents occurring in the derivation from S_0 up to S_k . A sequent $\Gamma \Rightarrow \Delta$ is saturated if it is not an instance of an initial sequent, and the following conditions are satisfied:

- (L_{\rightarrow}) If $x : A \rightarrow B$ occurs in $\downarrow \Gamma$, $x : B$ occurs in $\downarrow \Gamma$ or $x : A$ occurs in $\downarrow \Delta$;
- (R_{\rightarrow}) If $x : A \rightarrow B$ occurs in $\downarrow \Delta$, $x : A$ occurs in $\downarrow \Gamma$ and $x : B$ occurs in $\downarrow \Delta$,³
- (Ref) If y occurs in Γ , then $y \leq_x y$ occurs in Γ ;
- (Tr) If $y \leq_x z$ and $z \leq_x k$ occur in Γ , $y \leq_x k$ occur in Γ ;
- ($\text{L}_{>}$) If $x : A > B$ and $z \in W_x$ occur in $\downarrow \Gamma$, then either $z : A$ occurs in $\downarrow \Delta$ or $C_x^z(A, B)$ occurs in $\downarrow \Gamma$;
- ($\text{R}_{>}$) If $x : A > B$ occurs in $\downarrow \Delta$, then $z \in W_x$ and $z : A$ occur in $\downarrow \Gamma$, for some z and $C_x^z(A, B)$ occurs in Δ ;
- (LC) if $C_x^z(A, B)$ occurs in Γ , then either for some y $y \leq_x z$, $y \in W_x$, $y : \Box_x(A \rightarrow B)$ occur in $\downarrow \Gamma$, or for some w such that $z \neq w$, $z \leq_x w$ and $C_x^w(A, B)$ occur in $\downarrow \Gamma$;
- (RC) If $y \leq_x z$, $y \in W_x$, $z \in W_x$ occur in $\downarrow \Gamma$ and $C_x^z(A, B)$ occurs in $\downarrow \Delta$, then either $y : A$ or $y : \Box_x(A)$ occurs in $\downarrow \Delta$;
- (L_{\Box_x}) If $y : \Box_x A$ occurs in $\downarrow \Gamma$, $z \leq_x y$ and $z \in W_x$ occur in Γ , and $z : A$ occurs in $\downarrow \Gamma$;
- (R_{\Box_x}) If $y : \Box_x A$ occurs in $\downarrow \Delta$, then either for some z , $z \leq_x y$ occurs in Γ and $z : A$ occurs in $\downarrow \Delta$, or for some $w \neq y$, $y \leq_x w$ occurs in Γ and $w : \Box_x A$ occurs in $\downarrow \Delta$;
- (N) If x occurs in Γ , $y \in W_x$ occurs in Γ , for some y ;
- (T) If x occurs in Γ , $x \in W_x$ occurs in Γ ;
- (W) If $y \in W_x$ occurs in Γ , $x \leq_x y$ occurs in Γ ;
- (C) If $y \leq_x x$ and $y \in W_x$ occur in Γ , $y = x$ occurs in Γ ;
- (Ref₌) If x occurs in Γ , then $x = x$ occurs in Γ ;
- (Repl) If $y = x$ occurs in Γ , and if some formulas $\text{At}(y)$ occur in Γ , formulas $\text{At}(x)$ occur in Γ ;
- (Nes) If $y \in W_x$ and $z \in W_x$ occur in Γ , $y \leq_x z$ or $z \leq_x y$ occur in Γ .

³ The saturation conditions for the other propositional rules are standard [20].

In Example 1, the saturation condition (LC) blocks the application of the rule to formula $C_x^y(C, D)$, since $y \leq_x z$ and $C_x^z(C, D)$ occur in the antecedent. Intuitively, rule LC is not applied to a formula $C_x^y(C, D)$ if y has been generated by a previous application of LC to the same $C_x(C, D)$, possibly labelled with a different label, i.e., if for some z , formulas $y \leq_x z$ and $C_x^z(C, D)$ occur in a lower antecedents. A similar saturation condition is needed for $R\Box_x$.

Definition 6. *In root-first proof search for $\Rightarrow x_0 : A_0$, apply the following:*

1. Rules which do not introduce new labels are applied before rules which do introduce new labels;
2. A rule R cannot be applied to a sequent if the sequent already satisfies the saturation condition associated to R .

We need to show that every branch of a derivation starting with $\Rightarrow x_0 : A_0$ and built in accordance with the strategy is finite. Since labels can be attached only to the finitely many subformulas of formula A_0 , it suffices to prove that only a finite number of labels can occur in the branch. To this aim, we construct an acyclic graph with the labels occurring in the derivation, and show that the graph is finite: more precisely, that every node of the graph has a finite number of immediate successors, and that each branch of the graph is finite.

Definition 7. *Given a derivation branch as in Definition 5, let x, y be labels occurring in Γ . Let $k(x) = \min\{t \mid x \text{ occurs in } \Gamma_t\}$. We say that “ x generates y ”, in symbols xRy , if for some $t \geq k(x)$, $k(y) = t$ and $y \in W_x$ occurs in Γ_t .*

By inspection on the rules of $\mathbf{G3P}^{\mathbf{V}/\mathbf{N}/\mathbf{T}/\mathbf{W}/\mathbf{C}}$ and by definition of R we have that the relation R does not contain any cycles and forms a graph having at the root label x_0 , and that all the labels occurring in the derivation occur in the graph. The notion of conditional degree, needed to prove Lemma 2, corresponds to the level of nesting of the conditional operator $>$.

Definition 8. *The conditional degree of a formula A is defined as: $d(\perp) = d(p) = 0$; $d(A \circ B) = d(C_x^y(A, B)) = \max(d(A), d(B))$ for $\circ = \{\wedge, \vee, \rightarrow\}$; $d(\Box_k A) = d(A)$, and $d(A > B) = \max(d(A), d(B)) + 1$. For x a label in a derivation, $d(x) = \max\{d(C) \mid x : C \text{ occurs in } \downarrow \Gamma \cup \downarrow \Delta\}$.*

Lemma 1. *Every node in the graph generated by the relation R has a finite number of immediate successors.*

Proof. By definition, label y is generated from x if there exists a t such that y does not occur in Γ_s for any $s < t$, and $y \in W_x$ occurs in Γ_t . We need to prove that only a finite number of formulas $y \in W_x$ can be introduced from x .

Formulas $y \in W_x$ are introduced in root-first proof search by application of rules for semantic conditions, $R>$, LC or $R\Box_x$. In the first cases, $y \in W_x$ is introduced by \mathbf{N} or \mathbf{T} .⁴ By the saturation conditions, these rules can be applied

⁴ Observe that \mathbf{Repl} does not introduce new labels; however, it could introduce new links between the nodes of the graph. In the presence of \mathbf{Repl} the structure generated by R is a graph; otherwise, it is a tree.

at most once to a label x ; thus, they generate at most 2 new labels. If $y \in W_x$ is introduced by $R >$, the rule must have been applied to some $x : C > D$ occurring in Δ_{t-1} . By the saturation condition, rule $R >$ can be applied at most once to each formula $x : C > D$, and the number of such formulas linearly depends on $d(A_0)$, the degree of formula A_0 at the root of the tree. Similarly, rule $R\Box_x$ is applied to some $w : \Box_A$ in $\downarrow \Gamma_{t-1}$, generating a new $y \in W_x$. Formulas $w : \Box_A$ are introduced by RC and $R >$, which do not generate loops. The saturation condition ($R\Box_x$) ensures that no loops arise with formulas $L\Box_x$. Thus, only a finite number of labels can be introduced. In case $y \in W_x$ is introduced by LC the situation is more complex. As shown in Example 1, rule LC might interact with rule $L >$ generating a large number of new labels, however, thanks to the proof search strategy, their number is finite. We consider a case of loop more complex than the one in Example 1: suppose formulas $x : E_1 > F_1, \dots, x : E_k > F_k$ occur in the succedent of a sequent. Then, for some $z \in W_x$ in the antecedent, we can apply k times rule $L >$, generating k formulas $C_x^z(E_1, F_1), \dots, C_x^z(E_k, F_k)$. Then, rule LC can be applied to these formulas, generating k new labels $z_1 \in W_x, \dots, z_k \in W_x$, with $z_1 \leq z, \dots, z_k \leq_x z$. Moreover, the rule introduces in the antecedent formulas $z_1 : E_1, \dots, z_k : E_k$ and $z_1 : \Box_x(E_1 \rightarrow F_1), \dots, z_k : \Box_x(E_k \rightarrow F_k)$. Rule $L >$ can be applied to these labels, generating $k \cdot k$ new formulas:

$$\begin{array}{c} C_x^{z_1}(E_1, F_1), \dots, C_x^{z_1}(E_k, F_k) \\ \vdots \\ C_x^{z_k}(E_1, F_1), \dots, C_x^{z_k}(E_k, F_k) \end{array}$$

Application of LC to these formulas would in principle generate $k \cdot k$ new labels; however, the saturation condition (LC) blocks the application of the rule to all formulas. For $1 \leq i \leq k$ and $1 \leq j \leq k$, consider formula $C_x^{z_i}(E_j, F_j)$. It holds that formulas $z_i \leq_x z$ and $C_x^z(E_j, F_j)$ occur in lower antecedents/succedents, satisfying the saturation condition. Thus, for each $y \in W_x$, and for $k >$ -formulas occurring in the antecedent $k \cdot k$ new labels are generated. \square

Lemma 2. *Every branch in the graph generated by the relation R is finite.*

Proof. By induction on $d(x)$, for x a label in the graph. We show that the length of an arbitrary chain starting from x is bounded by the degree of the formula it labels. If $d(x) = 0$, the formulas labelled with x are all atomic or propositional formulas, and no formula $y \in W_x$ needs to be introduced. If $d(x) > 0$, there must be some formula $x : A > B$ occurring in $\downarrow \Gamma \cup \downarrow \Delta$. Thus, there is at least one chain of length greater than zero in the branch, and some label y such that xRy . Observe that y can occur only as label of formulas of smaller degree than the formulas labelled with x . More precisely, for all formulas $x : A > B$ with $d(A > B) \leq d(x)$ occurring in $\downarrow \Gamma \cup \downarrow \Delta$, it holds that for all formulas $y : \Box_x(A \rightarrow B)$, $d(y) < d(x)$, i.e., all labels introduced by combination of $R >$, $L >$, LC , RC and $R\Box_x$ are label of formulas with a smaller degree than formulas labelled with x . \square

It follows from Lemmas 1 and 2 that the acyclic graph is finite. Since the formulas occurring in a derivation are subformulas of the formula A_0 , and since the number of labels occurring in a derivation is finite, proof search terminates.

Theorem 4 (Termination). *Proof search in $\mathbf{G3P}^{\mathbf{V}/\mathbf{N}/\mathbf{T}/\mathbf{W}/\mathbf{C}}$ built in accordance with the proof search strategy for a sequent $\Rightarrow x_0 : A_0$ always comes to an end in a finite number of steps, and each sequent occurring as a leaf of the derivation tree is either an initial sequent or a saturated sequent.*

Termination of proof search allows to prove completeness by constructing a countermodel from a saturated sequent.

Theorem 5. *Let $\Gamma \Rightarrow \Delta$ be a saturated sequent in a $\mathbf{G3P}^{\mathbf{V}/\mathbf{N}/\mathbf{T}/\mathbf{W}/\mathbf{C}}$ derivation. There exists a finite countermodel $\mathcal{M}_{\mathcal{B}}$ satisfying all formulas in $\downarrow \Gamma$ and falsifying all formulas in $\downarrow \Delta$.*

Proof. The countermodel $\mathcal{M}_{\mathcal{B}}$ is constructed as follows: $W_{\mathcal{B}} = \{x \mid x \text{ occurs in } \downarrow \Gamma \cup \downarrow \Delta\}$; for all $x \in W_{\mathcal{B}}$, $W_x = \{y \mid y \in W_x \text{ occurs in } \Gamma\}$; $\leq_x = \{\langle y, z \rangle \mid y \leq_x z \text{ occurs in } \Gamma\}$; for p atomic, $\llbracket p \rrbracket = \{x \in W_{\mathcal{B}} \mid x : p \text{ occurs in } \Gamma\}$.

It is immediate to verify that the relation \leq_x satisfies the properties of reflexivity and transitivity; thus, $\mathcal{M}_{\mathcal{B}}$ is a model for \mathbf{PCL} . In the presence of \mathbf{N} , \mathbf{T} , \mathbf{W} , \mathbf{C} and \mathbf{Nes} , the saturation conditions associated to these rules ensure that the model $\mathcal{M}_{\mathcal{B}}$ is a model for the corresponding logic.⁵

Let ρ be the realization $\rho(x) = x$. We show that 1) if \mathcal{F} occurs in $\downarrow \Gamma$, $\mathcal{M}_{\mathcal{B}} \vDash_{\rho} \mathcal{F}$, and 2) if \mathcal{F} occurs in $\downarrow \Delta$, $\mathcal{M}_{\mathcal{B}} \not\vDash_{\rho} \mathcal{F}$.⁶ The two claims are proved by cases, and by induction on the weight of \mathcal{F} . If \mathcal{F} is a relational atom $y \in W_x$ or $y \leq_x z$ or a formula $x : p$, claim 1 (and claim 2) hold by definition of the model. The propositional cases and the cases of $\mathcal{F} = x : A > B$ and $\mathcal{F} = y : \Box_x A$ follow applying the inductive hypothesis. By way of example, we prove claim 2 for $\mathcal{F} = x : A > B$. Suppose that formula $x : A > B$ occurs in $\downarrow \Delta$. By the saturation condition associated to $\mathbf{R} >$, for some label z , $z \in W_x$ and $z : A$ occur in $\downarrow \Gamma$ and $C_x^z(A, B)$ occurs in Δ . Thus, by inductive hypothesis, $\mathcal{M}_{\mathcal{B}} \vDash_{\rho} z : A$ and $\mathcal{M}_{\mathcal{B}} \not\vDash_{\rho} C_x^z(A, B)$, thus by the truth condition for the conditional, $\mathcal{M}_{\mathcal{B}} \not\vDash_{\rho} x : A > B$. \square

As a consequence of Theorems 4 and 5 we have that any underivable sequent originates, in a finite number of steps, a saturated sequent which is used to define a countermodel. We therefore have:

Corollary 2 (Strong completeness). *Any sequent $\Gamma \Rightarrow \Delta$ is either derivable in $\mathbf{G3P}^{\mathbf{V}/\mathbf{N}/\mathbf{T}/\mathbf{W}/\mathbf{C}}$ or has a (finite) countermodel in the corresponding class of models.*

⁵ In case of centering it is convenient to define worlds as equivalence classes, to account for formulas $x = y$. Thus, $[x] = \{y \mid x = y \text{ occurs in } \downarrow \Gamma\}$ and $W^c = \{[x] \mid y \text{ occurs in } \downarrow \Gamma \cup \downarrow \Delta\}$. Centering follows from the saturation condition (C).

⁶ In case of centering, we also need to show that if $[x] \vDash_{\rho} A$ and $y \in [x]$, then $[y] \vDash_{\rho} A$, and that if $[x] \vDash_{\rho} A$ then $x : A$ occurs in $\downarrow \Gamma$. The proof follows from admissibility of \mathbf{Repl} in its generalized form [20].

Completeness of the proof systems is an obvious consequence:

Theorem 6 (Completeness). *If A is valid in one of the logics without uniformity and absoluteness, sequent $\Rightarrow x : A$ is derivable in the corresponding $\mathbf{G3P}^{\mathbb{V}/\mathbb{N}/\mathbb{T}/\mathbb{W}/\mathbb{C}}$ calculus.*

Completeness, along with termination, allow to define a decision procedure for the logics based on the labelled calculi. However, the resulting decision procedure would be of at least NEXPTIME complexity - thus, far from the known complexity bounds for the logics.⁷

6 Conclusion and Related Work

In this work, we introduced a family of uniform labelled calculi that capture in a modular way the conditional logic \mathbb{PCL} and its extensions, including Lewis' counterfactual systems. The calculi internalise the semantics of preferential models. This semantics, studied, among others, by Lewis and Burgess [5, 15], makes explicit reference to the comparative plausibility ordering among worlds, implicitly assumed in Lewis's sphere models.

Several labelled proof systems for conditional logics have been defined in the literature. A recent approach, based on the methodology of neighbourhood semantics of [18] and [19], is presented in [10] and gives a uniform family of labelled calculi for \mathbb{PCL} and its extensions. Neighbourhood semantics is a generalization of Lewis's sphere semantics; whereas the latter is adequate for \mathbb{V} , neighbourhood semantics covers also weaker conditional logics. When compared to labelled calculi based on neighbourhood semantics, the calculi $\mathbf{G3P}^*$ appear to be simpler: they use just one set of labels, whereas the calculi based on neighbourhood semantics need two sets of labels, for worlds and for neighbourhoods. It turns out that the preferential semantics already used to define labelled sequent calculi for Lewis's conditional logic \mathbb{VC} in [20] is sufficiently expressive to treat uniformly also the weaker extensions of \mathbb{PCL} .

Preferential models have already been used in [9] to define tableau calculi for \mathbb{PCL} and all its extensions, but with the important difference, with respect to our approach, of the addition of the *Limit Assumption*⁸ and the use of a strict relation of comparative similarity. Another semantically inspired approach can be found in [21], that presents a sequent calculus for system \mathbb{CK} and some of its extensions. These logics are the weakest conditional systems, and they are weaker than the logics considered in this article.

Internal calculi (i.e., proof systems in which sequents have a direct formula interpretation) for conditional logics have also been defined: in [2] nested sequent calculi for \mathbb{CK} and some of its extensions are developed, whereas in [1] a nested and optimal calculus for counterfactual logic \mathbb{V} can be found (refer to [11] for cases of extensions). Finally, display calculi for \mathbb{CK} have been introduced recently in [6].

⁷ Refer to [7] for complexity results for conditional logics.

⁸ The Limit Assumption states that there are no infinite descending \leq_x -chains.

With respect to the labelled proof systems $\mathbf{G3P}^*$, the internal calculi are less modular: they capture weaker logics, such as \mathbb{CK} , or subfamilies of the logics considered in this article, such as \mathbb{V} and its extensions. In particular, the definition of internal calculi for \mathbb{PCL} seems challenging: up to now, the only internal proof system known for it is the resolution calculus presented in [16]. The labelled approach treats in a modular way both \mathbb{PCL} and \mathbb{V} . The challenge of defining labelled calculi on the basis of preferential semantics lies in identifying a decomposition of the conditional operator in terms of simpler operators directly treatable by the sequent calculus rules. Here, this is done by introducing the indexed operator $\Box_x A$, similarly to [9,20], and the binary operator $C_x^z(A, B)$.

As underlined in Remark 1, by dropping the requirement of modularity it is possible to have simpler labelled calculi for sub-families of logics. We plan to define such calculi and analyse their termination in root-first proof search, to investigate the possibility of a better complexity bound for the corresponding logics. Furthermore, following [9], simpler rules could be defined also for $\mathbf{G3P}$ with the introduction of the Limit Assumption on preferential models. Finally, we plan to study the relationship of $\mathbf{G3P}^*$ with labelled sequent calculi for conditional logics based on neighbourhood models [10]. Via the correspondence between neighbourhood and preferential structures [3], we conjecture that the two families of calculi can be proved equivalent. It would be interesting to know which family of calculi allows for the better decision procedure in terms of complexity.

References

1. Alenda, R., Olivetti, N., Pozzato, G.L.: Nested sequent calculi for conditional logics. In: del Cerro, L.F., Herzog, A., Mengin, J. (eds.) JELIA 2012. LNCS (LNAI), vol. 7519, pp. 14–27. Springer, Heidelberg (2012). https://doi.org/10.1007/978-3-642-33353-8_2
2. Alenda, R., Olivetti, N., Pozzato, G.L.: Nested sequent calculi for normal conditional logics. *J. Logic Comput.* **26**(1), 7–50 (2013)
3. Alexandroff, P.: Diskrete Räume. *Mat.Sb. (NS)* **2**(3), 501–519 (1937)
4. Baltag, A., Smets, S.: A qualitative theory of dynamic interactive belief revision. *Log. Found. Game Decis. Theory (LOFT 7)* **3**, 9–58 (2008)
5. Burgess, J.P.: Quick completeness proofs for some logics of conditionals. *Notre Dame J. Formal Log.* **22**(1), 76–84 (1981)
6. Chen, J., Greco, G., Palmigiano, A., Tzimoulis, A.: Non normal logics: semantic analysis and proof theory. arXiv preprint [arXiv:1903.04868](https://arxiv.org/abs/1903.04868) (2019)
7. Friedman, N., Halpern, J.Y.: On the complexity of conditional logics. In: Doyle, J., Sandewall, E., Torasso, P. (eds.) Principles of knowledge Representation and Reasoning: Proceedings of the Fourth International Conference (KR 1994), pp. 202–213. Morgan Kaufmann Pub. (1994)
8. Galles, D., Pearl, J.: An axiomatic characterization of causal counterfactuals. *Found. Sci.* **3**(1), 151–182 (1998)
9. Giordano, L., Gliozzi, V., Olivetti, N., Schwind, C.: Tableau calculus for preference-based conditional logics: PCL and its extensions. *ACM Trans. Comput. Log.* **10**(3), 21 (2009)

10. Girlando, M.: On the proof theory of conditional logics. Ph.D. thesis, University of Helsinki (2019)
11. Girlando, M., Lellmann, B., Olivetti, N., Pozzato, G.L.: Standard sequent calculi for Lewis' logics of counterfactuals. In: Michael, L., Kakas, A. (eds.) JELIA 2016. LNCS (LNAI), vol. 10021, pp. 272–287. Springer, Cham (2016). https://doi.org/10.1007/978-3-319-48758-8_18
12. Girlando, M., Negri, S., Olivetti, N., Risch, V.: Conditional beliefs: from neighbourhood semantics to sequent calculus. *Rev. Symb. Log.* **11**, 1–44 (2018)
13. Grahne, G.: Updates and counterfactuals. *J. Log. Comput.* **8**(1), 87–117 (1998)
14. Kraus, S., Lehmann, D., Magidor, M.: Nonmonotonic reasoning, preferential models and cumulative logics. *Artif. Intell.* **44**(1–2), 167–207 (1990)
15. Lewis, D.K.: *Counterfactuals*. Blackwell, Oxford (1973)
16. Nalon, C., Pattinson, D.: A resolution-based calculus for preferential logics. In: Galmiche, D., Schulz, S., Sebastiani, R. (eds.) IJCAR 2018. LNCS (LNAI), vol. 10900, pp. 498–515. Springer, Cham (2018). https://doi.org/10.1007/978-3-319-94205-6_33
17. Negri, S.: Proof analysis in modal logic. *J. Philos. Log.* **34**(5–6), 507 (2005)
18. Negri, S.: Proof theory for non-normal modal logics: the neighbourhood formalism and basic results. *IFCoLog J. Log. Appl.* **4**, 1241–1286 (2017)
19. Negri, S., Olivetti, N.: A sequent calculus for preferential conditional logic based on neighbourhood semantics. In: De Nivelle, H. (ed.) TABLEAUX 2015. LNCS (LNAI), vol. 9323, pp. 115–134. Springer, Cham (2015). https://doi.org/10.1007/978-3-319-24312-2_9
20. Negri, S., Sbardolini, G.: Proof analysis for Lewis counterfactuals. *Rev. Symb. Log.* **9**(1), 44–75 (2016)
21. Poggiolesi, F.: Natural deduction calculi and sequent calculi for counterfactual logics. *Studia Logica* **104**(5), 1003–1036 (2016)
22. Weiss, Y.: *Frontiers of conditional logic*. Ph.D. thesis, The Graduate Center, City University of New York (2019)